

Banana bunchy top disease continues to be the main pest concern for banana growers across the state. The disease is caused by the *banana bunchy top virus* (BBTV). The banana aphid, *Pentalonia nigronervosa*, is the insect vector of the banana bunchy top virus. BBTV was first discovered on O‘ahu in 1989. Since then, the disease has been detected in the Kona, Hilo, and North Kohala areas of Hawai‘i and on Kaua‘i, Maui, and Moloka‘i. Continued spread of BBTV is a significant threat to Hawai‘i’s banana industry.

For more information about BBTV, please contact the Hawai‘i Department of Agriculture, your local Cooperative Extension office, or the Banana Bunchy Top website: www.ctahr.hawaii.edu/banana

Hawai‘i Department of Agriculture

O‘ahu 973-9538
 Maui 873-3558
 Hawai‘i 974-4140
 Kona 323-7594
 Kaua‘i 274-3069

University of Hawai‘i at Mānoa

College of Tropical Agriculture and Human Resources
 Cooperative Extension Service

Oahu

Agricultural Diagnostic Service
 Center 956-6706
 Honolulu 956-8139
 Pearl City Urban Garden Center ... 453-6050
 Wahiawa 622-4185
 Kaneohe 247-0421

Maui 244-3242

Moloka‘i 567-6929

Kaua‘i 274-3471

Hawai‘i

Hilo 981-5199
 Kamuela 887-6183
 Kona 322-4892

Produced by the Banana Action Group

UNIVERSITY OF HAWAI‘I AT MĀNOA

College of Tropical Agriculture and Human Resources
 (CTAHR)

UNIVERSITY OF HAWAI‘I AT HILO

College of Agriculture, Forestry and Natural Resource
 Management (CAFNR)

HAWAI‘I DEPARTMENT OF AGRICULTURE

Plant Industry Division

HAWAI‘I BANANA INDUSTRY ASSOCIATION

MAUI INVASIVE SPECIES COMMITTEE

U.S. DEPARTMENT OF AGRICULTURE

Agricultural Research Service
 Pacific Basin Agricultural Research Center

Published by the College of Tropical Agriculture and Human Resources (CTAHR) and issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Andrew G. Hashimoto, Director/Dean, Cooperative Extension Service/CTAHR, University of Hawai‘i at Mānoa, Honolulu, Hawai‘i 96822. An Equal Opportunity / Affirmative Action Institution providing programs and services to the people of Hawai‘i without regard to race, sex, age, religion, color, national origin, ancestry, disability, marital status, arrest and court record, sexual orientation, or status as a qualified veteran.

Production: Office of Communication Services, UH-CTAHR

Banana Bunchy Top

Symptoms and Control Methods



Clockwise from top left: adult female wingless aphid, adult winged aphid, and banana plants with severe bunchy top symptoms

PLEASE REMEMBER!

Do not transport banana plants
 into or between the
 islands of Hawai‘i.
 Transporting banana fruit
 is allowed.

Stop the spread!

With many small-acreage farms and urban backyard gardens in close proximity, BBTV suppression efforts in one location will not have a sustainable impact on the environment without a collaborative effort from everyone who grows bananas. It is imperative that BBTV prevention occurs in both banana-producing agricultural areas and in the community at large.

Insect Vector



Winged banana aphid Aphid colony on banana petiole

BBTV Best Management Practices

- 1) Disease Identification:** Bunchy top disease is diagnosed by symptom identification or by a lab test for the virus. If you suspect BBTV in your banana orchards, please contact the Hawai'i Department of Agriculture or your local Cooperative Extension office for more information. For a nominal fee, the UH-CTAHR Agricultural Diagnostic Service Center can test banana leaves for the banana bunchy top virus.
- 2) Insect Control:** Controlling aphids is essential to reducing the spread of BBTV. Aphid populations may increase during the spring or during warm, dry weather, and they can often be found on suckers. Please contact the Hawai'i Department of Agriculture, your local Cooperative Extension office, or the Banana Bunchy Top website <www.ctahr.hawaii.edu/banana> for a list of available aphicides.
- 3) Disease Prevention:** To minimize the spread of BBTV, all infected plants and their mats must be destroyed with an approved herbicide. Destroying infected plants with Roundup Ultra Max can help reduce the spread of BBTV. Fruits must be removed before treatment. Please read and follow the label instructions. Aphids on infected plants should be killed with an approved insecticide before injecting Roundup Ultra Max into a hole made at the base of the stem. Do not immediately cut the plants—the foliage is needed to move the herbicide throughout the plant. Allow injected plants to dry completely before moving them off your property.
- 4) Routine Monitoring:** Commercial and neighborhood banana plantings should be monitored for BBTV symptoms.
- 5) Alternative Host Management:** Control banana aphids on all known alternate hosts such as ginger, heliconia, and taro.
- 6) Fresh Start:** Backyard and new growers should avoid planting bananas until they can verify that their planting material is free of BBTV. Only BBTV-free plants should be used for planting.
- 7) Prevent Contamination:** Prevent the spread of BBTV by not transporting banana plants around your island or to other islands in the state.


Healthy



Leaves normal



Healthy petiole



No Morse code



No hooking



Healthy fruit

Infected



Leaves bunched up



Petiole with mottling



Morse code



Veins form green hooks



Stunted fruit