



# RONSTAR® G

## HERBICIDE

*Specimen Label*

- \* **NOT FOR USE ON HOME LAWNS.**
- \* **FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY.**
- \* **Selective Preemergent Herbicide for the Control of Annual Grasses and Broadleaf Weeds In Turf and woody Ornamental shrubs, vines, and trees.**

**ACTIVE INGREDIENT:**

Oxadiazon [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-  
 Δ-1, 3, 4-oxadiazolin-5-one] ..... 2.0%\*

**INERT INGREDIENTS:** ..... 98.0%

Total: ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**

### WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
 (If you do not understand the label find someone to explain it to you in detail).

EPA Reg. No. 432-886

EPA Est. No.

**IN CASE OF MEDICAL, ENVIRONMENTAL, OR  
 TRANSPORTATION EMERGENCIES OR INJURIES,  
 CALL 1-800-334-7577 (24 HOURS/DAY).**

**For Product Information,  
 Call Toll-Free: 1-800-331-2867**

NET CONTENTS:

**BACKED  
 by BAYER.**

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

## PRECAUTIONARY STATEMENTS

### WARNING

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes skin and eye irritation. Harmful if inhaled. Do not get on skin or on clothing. Avoid contact with eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants, waterproof gloves, a chemical-resistant apron when cleaning equipment, protective eyewear, chemical resistant headgear for overhead exposure and chemical resistant footwear plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. RONSTAR® should not be used in areas where irrigation or rainfall results in the direct or indirect (via ditches and canals) contamination of surface waters through dissolved runoff or erosion of soil particles to which oxadiazon is absorbed.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks and chemical-resistant headgear for overhead exposure.

Do not apply RONSTAR G through any type of irrigation system.

### STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food or feed by storage or disposal. Do not store in or around the home.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land fill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### GENERAL INFORMATION

RONSTAR G is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in turf and on a wide variety of woody ornamental shrubs, vines, and trees. RONSTAR G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Apply RONSTAR G with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

#### TURF

For use in established perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum\*, Tall Fescue, and Zoysia turf; such as, fairways, parks, golf courses, and lawns. Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results. Do not apply to wet turf. If rain is not expected shortly after application, irrigating turf thoroughly will increase effectiveness.

**\*Not currently registered in California.**

**PREEMERGENT WEED CONTROL**

WEEDS CONTROLLED	AMOUNT OF RONSTAR G		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass ( <i>Digitaria spp.</i> ) Goosegrass ( <i>Eleusine indica</i> ) *Field Sandbur ( <i>Cenchrus incertus</i> ) *Annual Sedge ( <i>Cyperus compressus</i> )	100 - 200 lbs.	2 ¼ - 4 ½ lbs.	Apply in late winter or early spring prior to seed germination. Application made after March or April may not be fully effective.
*Annual Bluegrass ( <i>Poa Annua spp.</i> ) <b>(See note below)</b>	100-200 lbs.	2 ¼ - 4½ lbs`	Apply in late summer to early fall prior to seed germination.
Virginia Buttonweed ( <i>Diodia virginiana</i> ) Pigweed ( <i>Amaranthus spp.</i> ) Florida Pusley ( <i>Richardia scabra</i> ) Oxalis ( <i>Oxalis spp.</i> ) Stinging Nettle ( <i>Urtica dioica</i> ) Carpetgrass ( <i>Axonopus offinis</i> )	200 lbs.	4 ½ lbs.	Apply in early spring prior to seed germination.

**NOTE:** In many areas, annual bluegrass exists as at least two biotypes or subspecies of *Poa annua*. The true annual biotype is *Poa annua* spp. *annua* which germinates every year from seed, while the biotype that behaves like a perennial is *Poa annua* spp. *reptans*. This perennial biotype can survive through the summer and established plants will not be controlled by this product. Consult the extension service or University weed specialists in your area for more information.

\*Not currently registered in California.

**FALL APPLICATION FOR SPRING WEED CONTROL IN COOL SEASON TURF\***

WEEDS CONTROLLED	AMOUNT OF RONSTAR G		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass, Goosegrass and other annual weeds	150 - 200 lbs.	3 3/8 - 4 1/2 lbs.	For use in Bluegrass, Bentgrass, Perennial Ryegrass and Tall Fescue turf. Make application in the fall after turf has become dormant. For late, fall applications, seeding into treated areas should be delayed until 4 months after greenup.

\*Not currently registered in California.

**ANNUAL WEED CONTROL DURING BERMUDAGRASS, ZOYSIAGRASS, SEASHORE PASPALUM, AND TROPIC LALO (*Paspalum Hieronymii*) ESTABLISHMENT FROM SPRIGS\***

WEEDS CONTROLLED	AMOUNT OF RONSTAR G		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass, Goosegrass, Annual Sedge, and other annual weeds	100 to 150 lbs.	2 ¼ to 3 ¾ lbs.	For Bermudagrass, make application immediately prior to or immediately after sprigging. For Seashore Paspalum, make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable Seashore Paspalum injury. Disturbance of the soil surface after RONSTAR G application may disrupt the herbicide barrier and reduce effectiveness. Irrigate immediately after application for best results.

For St. Augustinegrass and Bentgrass turf, do not exceed the rate of 150 pounds per acre (3 pounds active ingredient) of RONSTAR G.

Do not apply to Bentgrass turf mowed at less than ¼ inch.

RONSTAR G may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.

Avoid contact with flowers and shrubs except as recommended elsewhere on this label. Do not apply RONSTAR G to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment except as noted above. Do not apply to Dichondra or Centipedegrass. RONSTAR G should not be used on putting greens or tees because of the varying cultural practices used.

When making applications to Tropic Lalo, applications must be made 0 - 12 days after sprigging. Do not use in areas where endangered Hawaiian Goose or nene (*Branta sandvicensis*), Hawaiian duck (*Anas wyvilliana*), Hawaiian Stilt (*Himantopus mexicanus knudseni*), and Hawaiian Moorhen (*Gallinula chloropus sandvicensis*) may feed.

\*Not currently registered in California.

## WOODY ORNAMENTAL SHRUBS, VINES, TREES, AND CONIFER NURSERY APPLICATION

Apply RONSTAR G anytime during the year, prior to weed seed germination, for weed control. RONSTAR G can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of RONSTAR G will improve weed control activity. RONSTAR G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of RONSTAR G. Do not apply to wet foliage or under conditions in which granules will collect on leaves. Do not apply in greenhouses as plant injury may result from codistillation of RONSTARactive ingredient onto leaf surfaces in condensation.

### AMOUNT TO USE

Ground Application — Apply RONSTAR G broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 ¼ to 4 ½ pounds per 1,000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

\*Not currently registered in California for use in conifer nurseries.

ANNUAL BROADLEAF WEEDS	
Bittercress	( <i>Cardamine spp.</i> )
†Bristly Oxtongue	( <i>Picris echinoides</i> )
Carpetweed	( <i>Mollugo verticillata</i> )
†Cheeseweed	( <i>Malva parviflora</i> )
Common Groundsel	( <i>Senecio vulgaris</i> )
Common Purslane	( <i>Portulaca oleracea</i> )
*Evening Primrose	( <i>Oenothera spp.</i> )
†Fiddleneck	( <i>Amsinckia intermedia</i> )
†Fireweed	( <i>Epilobium angustifolium</i> )
*Galinsoga	( <i>Galinsoga spp.</i> )
†Garden Spurge	( <i>Euphorbia hirta</i> )
Golden Ragwort	( <i>Senecio aureus</i> )
Lambsquarters	( <i>Chenopodium album</i> )
**Liverwort	( <i>Marchantia spp.</i> )
†Niruri	( <i>Phyllanthus niruri</i> )
Pennsylvania Smartweed	( <i>Polygonum pensylvanicum</i> )
Petty Spurge	( <i>Euphorbia peplus</i> )
**Prostrate Spurge	( <i>Euphorbia supina</i> )
Redroot Pigweed	( <i>Amaranthus retroflexus</i> )
†Shepherdspurse	( <i>Capsella bursa-pastoris</i> )
†Sensitive Plant	( <i>Mimosa pudica</i> )
Sow Thistle	( <i>Sonchus oleraceus</i> )
Speedwell	( <i>Veronica spp</i> )
†Spiny Amaranth	( <i>Amaranthus spinosus</i> )
Spotted Catsear	( <i>Hypochaeris radicata</i> )
*Swinecress	( <i>Coronopus didymus</i> )
Yellow Woodsorrel	( <i>Oxalis stricta</i> )

ANNUAL GRASSES	
Annual Bluegrass	( <i>Poa annua</i> )
†Annual Sedge	( <i>Cyperus compressus</i> )
*Barnyardgrass	( <i>Echinochloa crus-galli</i> )
†Bristly Foxtail	( <i>Setaria verticillata</i> )
Crabgrass	( <i>Digitaria spp.</i> )
Fall Panicum	( <i>Panicum dichotomiflorum</i> )
Green Foxtail	( <i>Setaria viridis</i> )
Goosegrass	( <i>Eleusine indica</i> )
**Jungle Ricegrass	( <i>Echinochloa colonum</i> )
**Panama Paspalum	( <i>Paspalum fimbriaatum</i> )
†Ripgut Bromegrass	( <i>Bromus catharticus</i> )
†Wild Oats	( <i>Avena fatua</i> )

\*Use 150 to 200 pounds (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

\*\*Use 200 pounds (4.0 pounds of active ingredient) per acre for less susceptible weeds.

†Not currently registered in California.

## How To Calibrate Spreaders For Applications Of RONSTAR

Spreader Models	Ground Speed (MPH)	Rate RONSTAR G (Pounds Per Acre) Swath Width			(Feet)
		100	150	200	
CYCLONE B3-Pro	3	2.75	3.25	3.9	8
LELY HR*	3	3.25	3.75	4.25	21
	4	3.5	4.25	4.5	
	5	3.75	4.5	5.0	
LESCO ROTARY	3	C	D	E	7
SCOTT'S PRO R-8A	3	H	I+	J+	9
SPYKER 24-22	3	3.5	4.0	4.25	8
VICON	3	13	15	18	22
	4	15	18	22	
	5	16	21	25	

\* Settings based on use of small seed feeding with central spreader setting

NOTE: These settings are for initial calibration only. It is strongly recommended that your equipment be calibrated prior to actual application, using these settings as a starting point.

### CROP TOLERANCE

RONSTAR G is recommended for use on a wide variety of ornamental species and ground covers in container, field grown, and landscape planting situations. RONSTAR G may be used at the recommended rates on the following species.

#### ESTABLISHED ORNAMENTALS AND GROUND COVERS

<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>
Abutilon	<i>Abutilon pictum</i>	Banyan, Weeping	<i>Ficus benghalensis</i>
Acacia	<i>Acacia spp</i>	Barberry	<i>Berberis spp</i>
Agapanthus	<i>Agapanthus africanus</i>	Beafsteak plant	<i>Acalypha wilkesiana</i>
Allamanda, Common	<i>Allamanda cathartica</i>	Bearberry	<i>Arctostaphylos uva-ursi</i>
Alpine Willow	<i>Epilobium fleisheri</i>	Beech, European	<i>Fagus sylvatica</i>
Andromeda	<i>Pieris spp</i>	Birch, Paper	<i>Betula papyrifera</i>
†Anthuriums (Ozaki Red)		†Blanket Flower	Gaillardia aristata
Apache plume	<i>Fallugia paradoxa</i>	†Blazing Star	<i>Liatrix spicata</i>
Aralia	<i>Aralia spinosa</i>	Boxwood, Japanese	<i>Buxus microphylla</i>
Arborvitae	<i>Thuja spp</i>	Boxwood, Common	<i>Buxus sempervirens</i>
Ash, Green	<i>Fraxinus pennsylvanica</i>	Bougainvillea	<i>Bouganvillea spectabilis</i>
Azalea	<i>Rhododendron spp</i>	Brassaia, Dwarf	<i>Brassaia actinophylla</i>
†Baby's Breath	<i>Gypsophila paniculata</i>	Buddleia	<i>Buddleia davidii</i>
†Bald Cypress	<i>Takodium distichum</i>	†Bugleflower	<i>Ajuga reptans</i>
Bamboo, Feathery	<i>Bambusa vulgaris</i>	Callistemon	<i>Callistemon lanceolatus</i>

**ESTABLISHED ORNAMENTALS AND GROUND COVERS (con't)**

<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>
Camellia, Common, Japanese	<i>Camellia japonica</i>	Fountain Grass	<i>Pennisetum alopecuroides</i>
Camellia, Sasanqua	<i>Camellia sasanqua</i>	Gardenia	<i>Gardenia jasminoides</i>
Candytuft	<i>Iberis sempervirens</i>	Gazania, Trailing	<i>Gazania uniflora leucoleana</i>
Carissa	<i>Carissa macrocarpa</i>	Ginko	<i>Ginko biloba</i>
Carnation	<i>Dianthus caryophyllus</i>	†Gladiolas	
Ceanothus	<i>Ceanothus americanus</i>	†Golden Raintree	<i>Koelreuteria paniculata</i>
Cedar, Red	<i>Juniperus virginia</i>	Heath, Spring	<i>Erica carnea</i>
Cedar, Atlas	<i>Cedrus atlantica</i>	Heather, Scotch	<i>Calluna vulgaris</i>
Cistus	<i>Cistus purpureus</i>	Heliotropium	<i>Heliotropium arborescens</i>
Citrus, Ornamental*:		†Hemlock, Canada	<i>Tsuga canadensis</i>
Grapefruit	<i>Citrus paradisi</i>	Hibiscus	<i>Hibiscus spp</i>
Lemon	<i>Citrus limon</i>	Honeysuckle	<i>Lonicera spp</i>
Lime	<i>Citrus aurantifolia Bearss</i>	Holly	<i>Ilex spp</i>
Mandarin	<i>Citrus reticulata</i>	Ice Plant	<i>Carpobrotus edule</i>
Tangelo	<i>Citrus XTangelo Minneola</i>	Ilima	<i>Sida fallax</i>
Tangerine	<i>Citrus reticulata var. tangerina</i>	Ivy, Algerian	<i>Hedera canariensis</i>
Temple	<i>Citrus Xnobilis Temple</i>	Ivy, English	<i>Hedera helix</i>
Cuban Shaddock	<i>Citrus maxima</i>	Jasmine, Orange	<i>Murraya paniculata</i>
Valencia	<i>Citrus sinensis</i>	Jasmine, Primrose	<i>Jasminum mesnyi</i>
Washington Navel	<i>Citrus sinensis</i>	Jasmine, Star	<i>Trachelospermum jasminoides</i>
†Cleyera	<i>Cleyera laponica</i>	Jasmine, Showy	<i>Jasminum floridum</i>
†Coreopsis	<i>Coreopsis lanceolata</i>	Juniper	<i>Juniperus spp</i>
†Coromandel	<i>Asystasia gagentica</i>	Justica	<i>Justica brandegeana</i>
Cotoneaster	<i>Cotoneaster spp</i>	†Kwanzan Cherry	<i>Prunus serrulata</i>
†Crabapple*	<i>Malus spp</i>	†Lamb's Ear	<i>Stachys byzantina</i>
Crapemyrtle	<i>Lagerstroemia indica</i>	Lantana, Trailing	<i>Lantana montevidensis</i>
Creeping Phlox	<i>Phlox subulata</i>	Lapalapa	<i>Cheirodendron spp</i>
Chrysanthemum	<i>Chrysanthemum spp</i>	Larch, European	<i>Larix decidua</i>
Cypress	<i>Cupressus spp</i>	Laurel, English	<i>Prunus laurocerasus</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	†Leopards - Bane	<i>Doronicum cordatum</i>
†Dahlia	<i>Dahlia spp.</i>	Leucothoe, Coast	<i>Leucothoe axillaris</i>
Daphne, Fragrant	<i>Daphne odora</i>	Leucothoe, Drooping	<i>Leucothoe fontanesia (catesbaei)</i>
Deutzia, Slender	<i>Deutzia gracilis</i>	Lily of the Nile	<i>Agapanthus umbellatus</i>
†Dieffenbachia	<i>Dieffenbachia spp</i>	Lilac	<i>Syringa spp</i>
Dogwood, Common	<i>Cornus florida</i>	†Littleleaf Linden	<i>Tilia cordata</i>
Dogwood, Redosier	<i>Cornus stolonifera</i>	Little Pickles	<i>Othonna capensis</i>
Dogwood, Koussa	<i>Cornus koussa</i>	Loquat*	<i>Eriobotrya japonica</i>
Douglas fir	<i>Pseudotsuga menziesii</i>	Magnolia	<i>Magnolia spp</i>
ElmUlmus spp		Mahonia, Grape Holly	<i>Mahonia aquifolium</i>
Escallonia	<i>Escallonia spp</i>	Maple	<i>Acer spp</i>
Eucalyptus	<i>Eucalyptus spp</i>	Metallic Plant	<i>Hemigraphis colorata</i>
Euryops, Gray Leaved	<i>Euryops pectinatus</i>	†Mexican Orange	<i>Choisya ternata</i>
Euonymus, Spreading	<i>Euonymus kiautschovicus</i>	†Mock Orange	<i>Philadelphus virginialis</i>
Euonymus, Wintercreeper	<i>Euonymus fortunei</i>	Money Tree	<i>Pleomele marginata</i>
Falsecypress, Hinoki	<i>Chaemaecyparis obtusa</i>	Murraea	<i>Murraea exotica</i>
Falsecypress, Japanese	<i>Chaemaecyparis pisifera</i>	Murraea	<i>Murraea paniculata</i>
Flowering Quince	<i>Chaenomeles speciosa</i>	Myoporum	<i>Myoporum laetum</i>
Forsythia	<i>Forsythia intermedia</i>		



Russian Olive	<i>Eleagnus angustifolia</i>
Spruce	<i>Picea spp</i>
Trailing African Daisy	<i>Osteospermum fruticosum</i>
Wedelia	<i>Wedelia trilobata</i>
†Wisteria, Chinese	<i>Wisteria sinensis</i>
†Yew	<i>Taxus spp</i>

#### FOREST NURSERY APPLICATION

Seedlings of the following species can be treated with RONSTAR G Herbicide when at least 4 weeks old:

<b><u>COMMON NAME</u></b>	<b><u>SCIENTIFIC NAME</u></b>
Pine, Loblolly	<i>Pinus teada</i>
Pine, Eastern White	<i>Pinus strobus</i>
Pine, Slash	<i>Pinus caribaea</i>

#### SENSITIVE VARIETIES

Under certain growing conditions, the plant varieties listed below may be sensitive to RONSTAR G Herbicide. Growers should evaluate usage on a small scale prior to large scale applications.

<b><u>PLANT SPECIES</u></b>	<b><u>VARIETIES</u></b>
Andromeda ( <i>Pieris japonica</i> )	Snowdrift, Temple Bell
Azalea	Pink Champagne, Hershey, ( <i>Rhododendron spp</i> ), Leuchtenfuer, Memoire John Hearrens, Hino-Crimson, White Grandeur, Everest, Girard Roberta
Cotoneaster	Eichholz, Himalayan, Shangri-la ( <i>Cotoneaster spp</i> )
Crapemyrtle ( <i>Lagerstroemia indica</i> )	Petite Red Imp, Seminole Bush
Heath, Spring ( <i>Erica carnea</i> )	Mediterranean Pink
Hemlock, Canada ( <i>Tsuga canadensis</i> )	Pendula
Hibiscus ( <i>Hibiscus syriacus</i> )	Red Heart
Rhododendron	Daphnoides, Nova Zembla, ( <i>Rhododendron spp</i> ), Catawba Album
Spruce, Norway ( <i>Picea abies</i> )	Pumila
Spruce, White ( <i>Picea glauca</i> )	Densata
Viburnum ( <i>Viburnum spp</i> )	Davidii

\* For use only on plants that will not bear edible fruit within one year of RONSTAR G Herbicide application.

†Not currently registered in California.

## NOTES

## NOTES

## NOTES

---

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer Environmental Science. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Bayer AG, 2005

---

CHIPCO AND RONSTAR are registered trademarks of Bayer.



Bayer Environmental Science  
A Business Group of Bayer CropScience LP

Bayer Environmental Science  
2 T.W. Alexander Drive, Research Triangle Park, NC 27709  
[www.bayerprocentral.com](http://www.bayerprocentral.com)

CHIPCO RONSTAR G Herbicide (MASTER) Approved 7/2/98  
Address  
Ron G-SL-081303B