

---

REVIEW ARTICLE

---

**Prospects for the biological control of  
subterranean termites (Isoptera:  
Rhinotermitidae), with special reference  
to *Coptotermes formosanus***

**T.W. Culliney<sup>1\*</sup> and J.K. Grace<sup>2</sup>**

<sup>1</sup>Hawaii Department of Agriculture, Division of Plant Industry, 1428 South King Street, Honolulu, Hawaii 96814, USA; <sup>2</sup>Department of Plant and Environmental Protection Sciences, University of Hawaii, 3050 Maile Way, Honolulu, Hawaii 96822, USA

**Abstract**

Costs associated with subterranean termite damage and control are estimated to approach \$2 billion annually in the United States alone. The Formosan subterranean termite, *Coptotermes formosanus* Shiraki, is one of the more economically important subterranean species. In recent years, the shortcomings associated with conventional chemical control methods have prompted policymakers and scientists to evaluate the potential for biological control of subterranean termites (*C. formosanus* in particular), that is, to determine the potential for natural enemies – predators, parasitoids and pathogens – to suppress termite populations. Ants are the greatest predators of termites, and may have a considerable local impact on termite populations in some areas of the world. A few parasitoids of termites are known, but their potential for regulating termite populations seems negligible. Characteristics of the colony, such as a protected, underground location (and, for the *C. formosanus* nest, its modular and dispersed nature), are likely to limit the impact predators and parasitoids have on subterranean termites. Thus, there seems little potential for use of these agents for subterranean termite control. For various reasons, pathogenic organisms, such as viruses, bacteria, Protozoa, nematodes and most fungi, have shown little promise for use in biological termite control. However, research suggests that strains of two well-studied, endoparasitic fungi, *Beauveria bassiana* and *Metarhizium anisopliae*, when employed in baiting schemes, may offer the potential for at least some measure of subterranean termite control, although their successful use is compromised by a number of inherent biological limitations and logistical problems that have yet to be solved. Although not strictly in the realm of classical biological control, recent studies suggest that natural products, such as ant semiochemicals and fungal metabolites (siderophores), or their synthetic analogues, eventually might find a use in termite control programmes as repellents or insecticides in wood treatments or soil applications if stable formulations can be developed.

---

\*Fax: (808) 973 9533

E-mail: culliney@elele.peacesat.hawaii.edu

### Introduction

Of the more than 2700 termite species in the world, only a small minority is of any economic concern. For example, only 70–80 species cause significant damage to structures (Edwards & Mill, 1986). Although the poorer areas of the tropics and subtropics probably suffer more termite damage, it is in the developed countries, with their low action thresholds for urban pests, that the economic impact of termites has received the most attention (Su & Scheffrahn, 1998). For example, the estimated damage to wood and wood products caused by termites annually in the United States exceeds \$750 million, 95% of which is contributed by subterranean termites (family Rhinotermitidae) (Mauldin, 1986). Moreover, subterranean termites account for 80% of the approximately \$1.5 billion spent annually for termite control in the USA (Su, 1993).

The Formosan subterranean termite, *Coptotermes formosanus* Shiraki (Isoptera: Rhinotermitidae), is one of the more economically important termites in the warmer regions of the world (Su & Tamashiro, 1987). Not only are houses and other buildings attacked by this insect, but also telephone poles, boats, underground electrical and telephone cables, other finished goods (e.g. books, paper, fabric) and at least 48 species of living plants, including valuable crops and ornamentals (Lai *et al.*, 1983; Tamashiro *et al.*, 1987). In China, nesting by the termite has been reported to undermine dykes and cause flooding (Gao, 1987). Although limited in geographic range, in terms of damage potential, *C. formosanus* is the most serious termite pest in that country and Japan (Lin, 1987; Mori, 1987), as well as in the USA.

Control of *C. formosanus* and other subterranean species involves measures to prevent infestation of structures or remedial actions once prevention fails (Su & Tamashiro, 1987; Su & Scheffrahn, 1990; Yates *et al.*, 1997). Prevention of infestations may involve physical means (e.g. sand or steel-mesh barriers beneath foundations) or, more commonly, chemical controls (reviewed by Su & Scheffrahn, 1998). For decades, soil treatment with persistent insecticides has been widely used to protect structures from ground-nesting termites.

Whereas preventive measures can provide an effective, long-term barrier to termite penetration of structures, once an infestation has begun, remedial control of *C. formosanus* is extremely difficult due to the large population size, areal extent and foraging range of the subterranean colony. For example, in the USA, standard practice is to treat the soil beneath and around the structure to halt current invasion and prevent future attack. Usually, no attempt is made to find and destroy the colony, which may be situated anywhere within a volume of soil of hundreds of millions of cubic metres (Su & Tamashiro, 1987). Treatment of soil around the structure therefore kills only a small proportion of the colony, the majority of the population surviving to re-infest the structure, either by flying reproductives or by workers entering through an untreated or inadequately treated portion of soil (Su & Scheffrahn, 1990).

In view of these shortcomings, research efforts have focused on alternative means of termite management. For example, recent studies have demonstrated the efficacy of baiting schemes employing insect growth regulators in eliminating subterranean termite colonies (e.g. Su & Scheffrahn, 1993; Su, 1994; Su *et al.*, 1995; Grace *et al.*, 1996).

A need to evaluate the potential for biological control of subterranean termites (*C. formosanus* in particular) also has been recognized (HDOA, 1994; Grace, 1997; USDA, 1999). An earlier review of non-chemical methods of termite control in agriculture and forestry touched upon biological control (Logan *et al.*, 1990). The present paper reviews more specifically and comprehensively the available information concerning termite interaction with the biotic environment to assess the potential for control of subterranean termites through biological means.

### Biological control of termites

Biological control constitutes a more environmentally acceptable alternative to traditional chemical control measures. When successfully implemented, it can yield permanent, cost-effective management of pest populations with minimal environmental disturbance. Classical biological control programmes (i.e. the discovery, release, and colonization of natural enemies: arthropod predators and parasitoids, and pathogens) have proved successful against a variety of pests, and knowledge of the role natural enemies play in the ecology of these species has steadily accumulated (Huffaker & Messenger, 1976; DeBach & Rosen, 1991). However, data for termites are relatively scant. Although the influence of predation on termite distribution, abundance and population dynamics may be partially understood in a few cases (Wood & Johnson, 1986), the natural role of termite pathogens is much more obscure (Lee & Wood, 1971).

### Predators

Termites have a wide variety of predators, both opportunist and specialist, vertebrate and invertebrate. Attack occurs mainly upon alate reproductives or foraging workers outside the nest (Logan *et al.*, 1990), but a few predators attack termites within the nest (Sheppe, 1970). Among the vertebrates, fish, anurans, lizards, snakes, birds and mammals (including man) are known to take termite prey when available (Nutting, 1969). Insect and other arthropod groups reportedly preying opportunistically on termites include Scorpiones, Solifugae, Acari, Opiliones, Araneae, Chilopoda, Thysanura, Anisoptera, Blattaria, Mantodea, Phasmida, Gryllidae, Reduviidae, Neuroptera, Carabidae, Elateridae, Staphylinidae, Diptera and aculeate Hymenoptera (Pemberton, 1928a; Nutting, 1969; Sheppe, 1970; Phillipsen & Coppel, 1977; Wood & Sands, 1978; Johnson & Hagen, 1981; Pearce, 1997), although all orders of entomophagous insects probably contain at least some species that feed on termites (Deligne *et al.*, 1981).

Ants are the greatest enemies of termites in all regions of the world (Deligne *et al.*, 1981; Hölldobler & Wilson, 1990). A large percentage of ant species, including those from the two largest genera, *Pheidole* Westwood and *Camponotus* Mayr, are opportunistic predators of termites (Hölldobler & Wilson, 1990). Veeresh & Gubbaiah (1984) observed the longlegged ant *Anoplolepis longipes* (Jerdon) (Formicidae), a major agricultural pest in India, to prey facultatively upon termites of an unidentified species. Predation by doryline ants on colonies of *Cubitermes ugandensis* Fuller and *Bellicositermes natalensis* (Haviland) (both Termitidae) was witnessed by Williams (1959) and Bodot (1961), respectively. Yin (1994?) found *Dorylus orientalis* Westwood to be an important

natural enemy of *Odontotermes formosanus* (Shiraki) and *Macrotermes barneyi* Light (both Termitidae). Traniello (1981) observed foraging *Nasutitermes costalis* (Holmgren) (Termitidae) to be attacked by workers of the genera *Crematogaster* Lund, *Wasmannia* Forel, *Azteca* Forel, *Pachycondyla* F. Smith, and *Odontomachus* Latreille, as well as *Camponotus*. It is likely that ants account for a significant proportion of the probable near-100% mortality of swarming reproductives (Wood & Sands, 1978).

Termites may comprise a significant portion of the diet in some ant species. For example, Wilson & Oliver (1969) found that the greatest proportion (16.3%) of prey taken by fire ants *Solenopsis richteri* Forel (Formicidae) in cutover pine forest consisted of termites (*Reticulitermes* sp.). Laboratory experiments suggested that *Coptotermes formosanus* might be relatively more resistant to fire ant predation than *Reticulitermes* spp. owing to a higher proportion of soldiers in colonies of the former and their more aggressive response to disturbance (Wells & Henderson, 1993; cf. Cornelius & Grace, 1996, 1997).

Specialized arthropod predators of termites apparently are confined to a few ant species in the subfamilies Ponerinae and Myrmicinae (e.g. Arnold, 1915–1924; Wheeler, 1936; Weber, 1964; Levieux, 1966; Maschwitz & Mühlenberg, 1975; Ohiagu & Wood, 1976; Longhurst *et al.*, 1978, 1979a,b; Longhurst & Howse, 1979; Maschwitz & Schönegge, 1983; Lepage & Darlington, 1984), some of which may serve as important regulators of termite populations in natural ecosystems. For example, *Decamorium uelense* (Sanstchi) (Formicidae) consumed up to 632 *Microtermes* spp.  $m^{-2} y^{-1}$ , more than 74% of the *Microtermes* standing crop population and approximately one-eighth of annual production (Longhurst *et al.*, 1979b). *Megaponera foetens* (Fabricius) (Formicidae), a species noted for its organized raids on termites in the Ethiopian region, preyed upon *Macrotermes bellicosus* (Smeathman) (Termitidae) at a rate of 141 termites  $m^{-2} y^{-1}$ ; other species were consumed at the rate of 52 termites  $m^{-2} y^{-1}$  (Longhurst *et al.*, 1978). Predation on *M. bellicosus* was equivalent to slightly more than the annual standing crop and about one-seventh of the total annual production (Collins, 1981). *Megaponera foetens* may make repeated raids on a termite colony until it is destroyed (Sheppe, 1970).

#### Parasitoids

The main advantage to employing parasitoids for pest control derives from their high degree of host specificity, and parasitoids have been used successfully in biological control programmes worldwide. However, little information is available on parasitoids of termites, and none of it seems to have relevance to biological control. Certain species of the family Phoridae (Diptera) are parasitic on termites (e.g. Schmitz, 1938; Disney, 1988; Disney & Kistner, 1989, 1990), and might play a role in regulating termite densities (Disney, 1986). In other phorids, the maggot gaining entry into the host through a wound (Askew, 1971). Bridwell (1920) and Kühne & Becker (1974) reported *Scleroderma* spp. (Hymenoptera: Bethyilidae) parasitizing drywood termites (Kalotermitidae), but did not assess their effects on termite population dynamics.

In summary, while a few specialists may have a considerable local impact on termite numbers, most termite predators are opportunistic and exert a low level of

mortality that probably is compensated for by colony reproduction (Sheppe, 1970). Moreover, not all termite species are equally vulnerable to predation. Many of the termites attacked by the specialist predators build above-ground nests that are readily located and may be easier to exploit than are subterranean colonies. The well-hidden nest of *C. formosanus*, for example, which may be situated as deeply as 3 m below ground level, comprises an extensive gallery system, in which the main nest is connected by runways with several subnests (Zimmerman, 1948; Su & Tamashiro, 1987). The dispersed and modular nature of the *C. formosanus* colony, as well as its protected, underground location, may limit the impact predation has on this termite species. Similarly, evidence suggests that parasitoids may be of limited value in controlling populations of concealed (including subterranean) pests in general (Gross, 1991). Thus, there would seem to be little potential for use of predators or parasitoids to control *C. formosanus*.

#### Pathogens

Soil insects have been considered to be good candidates for microbial control (Burgess, 1981). The soil environment offers conditions highly favourable for sustaining infection and promoting epizootics: it is humid, protected from ultra-violet radiation and crowded. However, few confirmed pathogens have been isolated from termites (Lund, 1971; Zoberi & Grace, 1990a; Grace, 1994; Zoberi, 1995). Also, whereas many prospective termite pathogens have been tested in the laboratory, relatively few have been given field trials.

#### Viruses

Viral infection of termites has been little reported. Gibbs *et al.* (1970) isolated a virus infecting *Coptotermes lacteus* (Froggatt) (Rhinotermitidae), which was similar to acute paralysis virus of the honey bee *Apis mellifera* Linnaeus (Hymenoptera: Apidae). A nuclear polyhedrosis virus (NPV), obtained from caterpillars of *Spodoptera littoralis* Boisduval (Lepidoptera: Noctuidae), was infective to a laboratory colony of *Kalotermites flavicollis* (Fabricius) (Kalotermitidae) (Al Fazairy & Hassan, 1988). Termites died 2–10 days post infection, and, based only on the laboratory results, the authors suggested that control of *K. flavicollis* with NPV might be feasible. However, the potential of viruses for termite control has yet to be evaluated in the field (Al Fazairy & Hassan, 1993).

Accessibility of the pest to be controlled is the prime factor affecting the efficacy of viral pathogens. Insects that feed openly on the foliage of host plants are most easily treated, and the most promising results have been obtained against pests of this type (e.g. caterpillars, sawfly larvae) (Smith, 1967). Insects living in concealed habitats, such as the soil, are more difficult to infect. There are other drawbacks to the use of viruses to suppress pest populations: viruses kill their hosts slowly compared to other pathogens; environmental factors, such as rainfall and solar radiation, may reduce viral persistence in soil; mass production of viruses is hampered by the need for living hosts or tissue culture; lastly, viral formulations have had difficulty competing successfully, on the basis of performance and cost, with other pest control products, such as chemical insecticides or even other microbial agents (Klein, 1988; Fuxa, 1990).

### Bacteria

Little research has focused on bacterial infections of termites. Exposure of laboratory colonies of the subterranean species *Reticulitermes flavipes* (Kollar) and *R. hesperus* Banks (Rhinotermitidae) to a mixture of soluble endotoxin, spores and inclusion bodies of *Bacillus thuringiensis* (Berliner) (*Bt*, a bacterium commonly used as a microbial insecticide for control of pest Lepidoptera, Coleoptera and Diptera) resulted in greater than 95% mortality after six days (Smythe & Coppel, 1965). Workers of *Microcerotermes championi* (Snyder) (Termitidae) and *Heterotermes indicola* (Wasmann) (Rhinotermitidae) suffered 100% mortality within 13 days of exposure to two local strains of *B. thuringiensis* in laboratory tests (Khan *et al.*, 1977a). In follow-up studies employing a commercial preparation of *Bt* (Thuricide-HP concentrate), Khan *et al.* (1978, 1985) again found *H. indicola*, *M. championi* and another termite species, *Bifiditermes beesoni* (Gardner) (Kalotermitidae), to be highly susceptible to infection, exhibiting 100% mortality within six days of exposure.

Laboratory colonies of *M. championi*, *H. indicola* and *B. beesoni* exposed to suspensions of the spore-forming bacterium *Serratia marcescens* Bizio succumbed completely 7–13 days following infection (Khan *et al.*, 1977b). Khan *et al.* (1992) tested the pathogenicity of *Pseudomonas aeruginosa* (Schroeter) against *M. championi*, *H. indicola* and *Coptotermes heimi* (Wasmann) (Rhinotermitidae) in the laboratory. Termite mortality ranged from 25–52% seven days post-inoculation to 84–100% 25 days post-inoculation. The authors concluded from the results that *P. aeruginosa* is 'fairly' pathogenic to the three termite species, although the bacterium's potential as a biological control agent is limited by its occasional status as a plant pathogen.

Despite favourable laboratory results, the potential of bacteria to reduce termite populations has not been demonstrated under field conditions. *Bacillus thuringiensis*, in particular, has limited potential for soil insect control because of its poor survival in soil (Burgess, 1981). Moreover, the insects that *B. thuringiensis* strains typically kill in the field are not soil-dwelling, but epigeal or aquatic (Martin, 1994). However, recent work incorporating recombinant DNA and microencapsulation technologies shows promise for application to termite control strategies employing *Bt* strains that express termite-specific endotoxin (Grace & Ewart, 1996).

### Protozoa

Of the four groups of Protozoa containing species parasitic to insects, the Phylum Microspora includes species that are potentially most useful in applied insect control (Henry, 1990). However, the amount of research pertaining to biological control of termites using these agents is negligible. Jafri *et al.* (1976) found microsporidians in the body cavity and proventriculus of *M. championi* collected from the roots of *Saccharum munja* Roxburgh (Poaceae). The organisms apparently are ingested with the food. After passing into the midgut, they attack fat body tissues and cause death. Desportes (1963) described a gregarine (Phylum Apicomplexa) from the haemocoel of the dampwood termite *Zootermopsis nevadensis* (Hagen) (Termitidae). No assessment of the microbes' potential for termite control was given.

Although Protozoa may be important agents in the natural control of many insects, they have not been used as soil-applied microbial insecticides because they tend to be slow-acting and vulnerable to changes in environmental conditions (Klein, 1988; Henry, 1990).

### Nematodes

The roundworms of the Phylum Nematoda play an important role in the natural control of many insect populations. They have been found parasitizing species in the orders Hemiptera, Diptera, Hymenoptera, Lepidoptera, Orthoptera, Coleoptera, Thysanoptera, Siphonaptera, as well as Isoptera (Nickle & Welch, 1984).

Four families of nematodes – the Mermithidae, Allantonematidae, Steinernematidae and Heterorhabditidae – have shown promise for use in insect control programmes (Popiel & Hominick, 1992), with most research focusing on species in the latter two (Kaya & Gaugler, 1993). These entomopathogenic nematodes (so-named because they are vectors of pathogenic bacteria of the genus *Xenorhabdus* Poinar) have a number of characteristics that make them especially suitable for biological control and for commercial production as microbial insecticides: a broad host range, especially among soil-dwelling pests; exemption from registration in the USA; ease of production, storage and application; a high degree of safety to vertebrates, plants and other non-target organisms; and amenability to genetic selection (Kaya & Gaugler, 1993; Kaya *et al.*, 1993).

Few studies have addressed the potential for nematodes to control termites. In an early report, Merrill & Ford (1916) found 77% of colonies, and up to 100% of individuals, of *Reticulitermes lucifugus* (Rossi) (Rhinotermitidae) to be parasitized by *Mikolletzkya aerivora* (Cobb) (Diplogastridae) in field samples. Laboratory exposure of termites to nematodes resulted in successful parasitization (47% at 4 days) and 100% mortality of infected individuals at 12 days. However, no field trials were undertaken. Fujii (1975) obtained 96% mortality in *C. formosanus* within seven days of exposure to infective-stage *Steinernema carpocapsae* (Weiser) (Steinernematidae) in laboratory experiments. Mortality exceeding 95% was recorded by Georgis *et al.* (1982) for both *Zootermopsis* sp. and *Reticulitermes* sp. within three days after laboratory exposure to *S. carpocapsae*; termites also were found to carry infection back to their colonies. Laumond *et al.* (1979) and Trudeau (1989) also reported high rates of infection of *Nasutitermes costalis* and *Reticulitermes flavipes*, respectively, with *S. carpocapsae* under laboratory conditions.

Results of field studies evaluating the efficacy of nematodes in termite control generally have been disappointing. In an early, informal test, Pemberton (1928b) introduced unidentified nematodes from an Indonesian species of *Termes* Linnaeus (Termitidae) into soil around a nest of *C. formosanus*. No evidence of subsequent parasitization was detected. In field trials, in which concentrations of *S. carpocapsae* as high as 10 million individuals m<sup>-2</sup> were applied to soil beneath wooden baits, Epsky & Capinera (1988) found evidence that *Reticulitermes tibialis* (Banks) (Rhinotermitidae) avoided contact with nematodes and exploited gaps in coverage in nematode-inoculated areas to attack the bait. The field results contrasted with those from their laboratory experiments, in which no avoidance of nematode-infested areas was found.

In a similar study using *Heterorhabditis heliothidis* (Khan, Brooks & Hirschmann) (Heterorhabditidae) as well as *S. carpocapsae* against *Reticulitermes* spp., nematodes applied to plots were not found to reduce feeding damage to wooden bait blocks, and showed no effectiveness in controlling or eliminating the termites (Mauldin & Beal, 1989).

Perhaps the only example of apparently successful biological control of a termite with nematodes involved not a subterranean, but a drywood species, *Glyptotermes dilatatus* (Bugnion & Popoff) (Kalotermitidae), a pest of tea, *Camellia sinensis* (L.) Kuntze (Theaceae). Danthanarayana & Vitarana (1987) showed that a single application of nematodes (*Heterorhabditis* sp.; 4000–8000 ml<sup>-1</sup> of suspension) to galleries in tea plants eliminated termite colonies within two to three months. Evidence suggested that nematode populations were self-perpetuating in the field, even under extreme environmental conditions. However, at a cost of US\$2.39 per 1000 infested bushes, this time-consuming and labour-intensive method is unlikely to be appropriate for use in less valuable crops or in forestry.

Although it may be feasible in limited cases to control crop-infesting termites with nematodes, in general, the utility of nematodes in biological control programmes is compromised by a number of factors that influence their effectiveness. These include soil physical and chemical properties (e.g. moisture, temperature, pore size, compaction, oxygen and carbon dioxide levels, pH, salinity and presence of synthetic chemicals) and biotic factors, such as competitive or predatory interactions with other soil organisms, limited motility and termite behaviours (e.g. isolation of infected individuals from the rest of the colony) that limit the spread of infection (Gaugler, 1988; Kaya, 1990; Popiel & Hominick, 1992).

Protection of structures from subterranean termites involves other considerations. In particular, the low economic threshold for termite damage to homes and other buildings, the immense numbers of nematodes required to kill termites, the presence of secondary reproductives that can rapidly re-establish the old colony or found new ones, and termite avoidance of nematode barriers all combine to limit the use of nematodes for termite control (Klein, 1990).

Although field infections often result from nematode treatments, the impact on target pest populations most often has been modest or negligible, even when billions of nematodes ha<sup>-1</sup> were applied (Kaya & Gaugler, 1993). Thus, the potential for use of nematodes in biological control may be limited to cases in which chemical insecticides are not practical or appropriate (Nickle & Welch, 1984).

### Fungi

Much research interest, especially in recent years, has focused on the use of fungal agents for pest control (Ferron, 1978). At least 22 species of fungi are obligate ectoparasites of termites (Blackwell & Rossi, 1986), although their biological control potential apparently has not been evaluated. Several of the fungi associated with termites appear to be at least facultatively pathogenic (e.g. Thaxter, 1895, 1920; Tate, 1927, 1928; Colla, 1928; Kevorkian, 1937; Stifler, 1941; Altson, 1947; Williams, 1959; Buchli, 1960; Beal & Kais, 1962; Lund & Engelhardt, 1962; Steinhaus & Marsh, 1962; Becker & Kerner-Gang, 1964; Toumanoff, 1965; Yendol & Paschke, 1965; Smythe & Coppel, 1966; Sannasi, 1968; Gouger & Kimbrough, 1969; Kimbrough & Gouger, 1970;

Kimbrough *et al.*, 1972; Khan & Kimbrough, 1974a,b; Rossi, 1974; Blackwell & Kimbrough, 1976, 1978; Blackwell, 1980; Blackwell & Gilbertson, 1981, 1984; Hänel, 1982b; Ko *et al.*, 1982; Preston *et al.*, 1982; Zoberi & Grace, 1990b; Yoshimura *et al.*, 1992; Ochiel, 1995; Ochiel *et al.*, 1997; Sajap *et al.*, 1997; Yoshimura & Takahashi, 1998). It is the endoparasitic fungi, however, that have perhaps the greater potential for biological control of termites, and these have received more attention by researchers.

Only two fungal pathogens, *Beauveria bassiana* (Balsamo) Vuillemin and *Metarhizium anisopliae* (Metschnikoff) Sorokin (Hyphomycetales), have been extensively evaluated for termite control. Rosengaus & Traniello (1997) tested the pathogenicity of *M. anisopliae* on laboratory colonies of *Zootermopsis angusticollis* (Hagen) (Termopsidae). Mortality due to infection was dosage-dependent, LT<sub>50</sub> (median survival time) values ranging from 2.7 days at a concentration of  $7 \times 10^7$  conidia (spores) ml<sup>-1</sup> to more than 10 days at  $7 \times 10^1$  conidia ml<sup>-1</sup>. Termites directly exposed to the highest concentration were capable of infecting unexposed, healthy nest mates. In a series of laboratory studies, Hänel (1981, 1982a,b) investigated the pathogenicity and biological control potential of *M. anisopliae* on the mound-building species *Nasutitermes exitiosus* (Hill) (Termitidae). There was a clear correlation between conidial concentration and percentage mortality, average mortality exceeding 95% within 11 days at the highest dosages. Termites were observed to segregate cadavers from the rest of the colony, but, in so doing, came into contact with infective conidia.

Subsequent field experiments did not confirm the potential of *M. anisopliae* to give permanent control of *N. exitiosus* colonies (Hänel & Watson, 1983). In five out of seven mounds treated with conidial preparations, termite population densities were significantly reduced. Also, termites exposed to conidia at feeding sites were seen to spread infection back to their colonies, which persisted in some for at least six weeks. Again, densities in the affected colonies were significantly decreased. However, although the fungus persisted for as long as 15 weeks in termite mounds, its incidence declined progressively, as did termite mortality, which was high initially, but decreased over time. The relevance of these results to the control of subterranean termites is unclear.

Several laboratory studies have examined effects of *M. anisopliae* and *B. bassiana* on subterranean termite populations. Results have varied with the termite species and strain of fungus tested, and the concentration of inoculum and its mode of delivery to termite populations. Leong (1966) demonstrated the high pathogenicity of *M. anisopliae* to *C. formosanus*. With short exposures (e.g. 5–35 min), mortality exceeded 86%, and termites succumbed within 3–6 days post treatment. Exposure over 40 min produced 100% mortality; longer exposure times caused death within 24 h. The fungus sporulated only sparsely under conditions of complete darkness, which might limit its utility for biological control of subterranean termites, although pathogenicity was unchanged between dark and light conditions. No significant difference between the pathogenicities of *B. bassiana* and *M. anisopliae* was found. In another study, lengthy exposure (4 h) to sporulating cultures of *B. bassiana* also led to the rapid death of *R. flavipes*, generally in 24–36 h (Bao & Yendol, 1971). Toumanoff & Rombaut (1965) reported 100% mortality in *Reticulitermes*

*santonensis* (de Feytaud) (Rhinotermitidae) within 5–6 days of exposure to *B. bassiana* and *M. anisopliae*. Sajap & Kaur (1990) also showed *Coptotermes curvignathus* (Holmgren) (Rhinotermitidae) to be highly susceptible to infection by *M. anisopliae*. Termites died within 36–48 h of exposure to conidia. Another study of *M. anisopliae*, in which termites exposed to conidia were introduced into laboratory colonies, reported 100% mortality in the test colonies within eight days, even when the termite vectors carried low conidial loads (Zoberi, 1995). Spread of the disease through colonies was complete, even though some of the fungus-killed termites were buried by nest mates, impairing growth of the fungus and preventing dispersal of conidia. Results suggested that epizootics within natural subterranean colonies might be initiated by a sufficient number of vector termites contaminated with large doses of inoculum introduced directly into the soil.

Kramm & West (1982) found that conidia of *B. bassiana*, *M. anisopliae* and *Gliocladium virens* Miller, Giddens & Foster remained infective after passage through the intestines of subterranean termites (*Reticulitermes* sp.), a significant finding since mutual grooming, trophallaxis and cannibalism should provide efficient means for spreading the pathogens throughout the colony. Again, strains of *B. bassiana* and *M. anisopliae* were about equally effective, producing complete mortality within 2–5 days. Similarly, Wells *et al.* (1995) concluded that one isolate each of *B. bassiana* and *M. anisopliae*, of all strains tested, showed the greatest potential for control of *C. formosanus* populations, based on LD<sub>50</sub> (median lethal dose; as conidia per insect), time to death, and conidial production.

However, Lai *et al.* (1982), using other strains of the fungi, found that *M. anisopliae* was more pathogenic than *B. bassiana* in conidial suspensions topically applied to *C. formosanus* colonies. A large increase in conidial concentration (at least 10-fold) was required to increase significantly the rate and extent of infection in the colonies. Grace (1991) also found *M. anisopliae* to cause greater and more rapid mortality than *B. bassiana* in *C. formosanus* at low conidial concentrations. In their evaluation of both pathogens, Jones *et al.* (1996) found the *M. anisopliae* strains to be more virulent, in general, than those of *B. bassiana*. However, although the *M. anisopliae* strains showed lower LT<sub>50</sub> values, the authors concluded that one strain of *B. bassiana* possessed greater potential as a remedial control for *C. formosanus*, based, in sum, on its lower LC<sub>50</sub> value (median lethal concentration), moderately low LT<sub>50</sub> and transmissibility and survivorship qualities.

Rath & Tidbury (1996) found that *Coptotermes acinaciformis* (Froggatt) (Rhinotermitidae) and *N. exitiosus* were equally susceptible to direct conidial applications of both an Australian and American strain of *M. anisopliae*. They concluded that virulence of *M. anisopliae* is not an essential strain selection criterion, as all isolates are apparently virulent. However, Milner (1992) found that, in general, *C. acinaciformis* and *C. frenchi* (Hill) (Rhinotermitidae) were more susceptible to *M. anisopliae* than *N. exitiosus*. Subsequently, an isolate of the fungus, which was highly pathogenic to all three termite species in the laboratory, reportedly showed some potential to control termite populations in field trials (Milner *et al.*, 1996), although few details were provided.

Other work has shown *M. anisopliae* and *B. bassiana* to be ineffective in controlling subterranean termites in the field.

In experiments in which conidia of the two fungi were introduced directly into galleries and onto termites subsequently returned to the nest, Lai (1977) failed to initiate an epizootic in colonies of *C. formosanus*, even though the concentration used was sufficient to kill termites within 3–7 days in the laboratory. The negative results were attributed to poor germination of conidia in the nest and elimination of conidia from treated termites through grooming or secretion of a fungistatic agent. Others also have been unable to demonstrate field efficacy of the pathogens despite successful laboratory trials (Suzuki, 1991, 1996; Gitonga, 1996). To date, injection of large quantities of conidia directly into the termite nest has had the greatest success in field studies (Fernandes, 1991; Milner *et al.*, 1996).

*Beauveria bassiana* and *M. anisopliae* may exhibit their greatest potential for subterranean termite control in their use in baiting systems, whereby the termites themselves pick up and introduce infective agents into the colony. Laboratory studies have begun to identify both the promise and problems associated with such techniques. For example, Preston *et al.* (1982) found that baits composed of *M. anisopliae* conidia and culture medium (containing a mixture of conidia, mycelia and metabolites), incorporated in different combinations into paraffin-woodmeal composite blocks, exhibited uniform repellency to termites (*Reticulitermes* sp.), but induced differential mortality. Blocks containing conidia alone produced no detectable mortality, whereas the highest mortalities (but least feeding damage) were associated with blocks containing medium, which remained effective as long as one month. However, the fungus could not be isolated from cadavers. This result, combined with the repellency of the baits, casts some doubt on the system's practical capacity to infect a natural colony. Other studies have reported the conidia (Rath & Tidbury, 1996) and toxins (destruxins; Wahlman & Davidson, 1993; Grace, 1995) produced by *M. anisopliae* to be repellent to termites.

Recent work with a *B. bassiana* strain isolated from *R. flavipes*, and pathogenic to that species (Zoberi & Grace, 1990a) and to *C. formosanus* (Grace, 1991), showed that, although *R. flavipes* workers isolated fungus-killed individuals, there was no avoidance of mycelia or conidia (Zoberi & Grace, 1990a). In a later study, Grace & Zoberi (1992) found that living *R. flavipes* workers exposed to sporulating *B. bassiana* cultures effectively spread infection to unexposed nest mates, whereas introduction of fungus-killed workers did not result in sufficient spore transfer or mycelial growth to cause significant mortality. However, the level of mortality achieved in the laboratory was not considered sufficient to control a termite infestation in the field. Similarly, Kramm *et al.* (1982) showed that active *Reticulitermes* sp. workers exposed to whole cultures of *M. anisopliae* transferred the pathogen to previously healthy termites through grooming. However, termites killed by the fungus were avoided by healthy individuals, and were less effective in spreading the disease to nest mates.

Other laboratory studies of baits have assessed the infectivity of sporulating strains of *B. bassiana* and *M. anisopliae* to *C. formosanus*. In preliminary work, Grace (1993) reported that termites exposed to baits accumulated spore loads (600,000–8 million conidia per individual) that greatly exceeded previously established lethal concentrations (LC<sub>95</sub>; 490–20,000 conidia per individual), and were capable of transmitting the pathogens to other colony members. Delate

*et al.* (1995) achieved complete mortality of termites within 15 days with isolates of both fungi. However, in this case, neither avoidance of fungi by termites nor isolation of infected individuals from the rest of the colony was detected. Results suggested that fungal bait stations might be useful in termite control by providing a continuous and non-repellent source of sporulating cultures for foraging termites to contact, although a self-perpetuating infection within the experimental colonies was not demonstrated.

In summary, of all pathogens tested, the fungi would seem to offer the most potential for at least some measure of termite control. However, the use of fungi in control programmes is compromised by inherent biological limitations and logistical problems. Fungi have a slow mode of action and require high levels of relative humidity; there is also the need for large quantities of infective conidia to contact the target pest population in order to yield an acceptable level of control (McCoy, 1990). It has also been suggested that, because so few of the many strains evaluated for termite control were effective, even under favourable conditions, termites may have evolved a degree of resistance to fungal pathogens (Burgess, 1981). However, there is as yet no good evidence to support this speculation.

Perhaps because of the many difficulties involved with their use, there are, at present, few commercial fungal preparations available for pest control in the USA. Further development for biological pest control of *B. bassiana* and *M. anisopliae*, in particular, continues to be hampered by a lack of cost-effective methods for mass production (Federici, 1990).

Fungi may yet find their most effective use in the form of baits, and research in this area is progressing (e.g. Delate *et al.*, 1995; Jones *et al.*, 1996; Staples & Milner, 1996). Effective baiting schemes for termite control require delivery of sufficient spore inoculum to active termites and subsequent transfer from them to nest mates without stimulating colony defensive behaviours. This will entail both improvement of field techniques for dispensing conidia and identification of new fungal species and strains capable of initiating and sustaining an epizootic once they are introduced into the nest.

### Conclusions

Although laboratory studies have hinted at its potential, classical biological control of termites has not yet been convincingly demonstrated. The limited number of field trials that have been attempted largely have failed to establish the permanent, self-sustaining populations of control agents necessary to reduce termite densities below economically damaging levels. This lack of success has resulted from a combination of factors. Intrinsic limitations of natural enemies, such as narrow environmental tolerances, antagonistic interactions with other organisms, limited mobility, or a low level of virulence, diminish their effectiveness as agents of control, whereas natural defensive behaviours of the termite prey, such as avoidance of areas of predator, parasite or pathogen concentration, detection and isolation of infected nest mates, and, in the case of subterranean termites, construction of a cryptic, complex, and largely inaccessible nest, reduce the impact of predation and infection. As Logan *et al.* (1990, p. 324) have remarked, 'Manipulation of predators, either local or introduced, will have little effect if only foraging worker termites are killed, while the possibility of manipulating predators which

seek out the king and queen seems remote. Many microorganisms are pathogenic to termites, but few have received more than cursory attention. This, combined with the termites' behavioural mechanisms for combatting infection of the colony, means that successful and widely applicable microbial control is a long way off.' An additional obstacle to progress in microbial termite control relates to the constraints that evolving environmental protection laws place on research, making it increasingly difficult for researchers to import insect pathogens.

A thorough assessment of biological termite control should address the possibility of unintended results. Any pest control programme poses some threat to the environment. Natural enemies introduced for the biological control of specific pests have had negative impacts on populations of nontarget organisms either directly, through predation, parasitism, infection or competition, or indirectly, in their influence on seemingly unconnected species (Howarth, 1991; Secord & Kareiva, 1996). Introduced subterranean termite enemies conceivably might adversely affect nontarget termite species or have secondary effects on other organisms coming into contact with dead or dying termites. To minimize such risks, governments of developed countries, such as Australia and the United States, in recent years have adopted strict quarantine regulations governing the importation of exotic organisms. In the USA, for example, the Department of Agriculture (USDA) and Environmental Protection Agency (EPA) oversee the introduction, host or prey specificity testing and release of biological control agents (Pimentel *et al.*, 1984; Secord & Kareiva, 1996). The Environmental Protection Agency is the lead agency responsible for registering and regulating microbial pesticides (Grace, 1994). The testing protocols and other quarantine procedures developed by these agencies are the best compromise between the competing demands of pest control and environmental protection.

However disappointing near-term prospects for successful classical biological termite control would seem to be, research has begun to test promising new parabiological methods to prevent structural infestation by subterranean termites. For example, recent studies have examined the potential for use of ant semiochemicals as termite repellents and toxicants. In laboratory tests conducted by Cornelius & Grace (1994), *C. formosanus* workers avoided contact with filter paper disks treated with extracts of the ant *Ochetellus glaber* (Mayr) (Formicidae); no evidence of termite habituation to the extracts was detected. Extract-treated sand barriers deterred tunnelling completely for 2-4 days at the higher extract concentrations tested, although partial penetration was seen in succeeding days. However, none of these barriers was completely penetrated, even after ten days. Further work showed sand treated with *O. glaber* extracts to be toxic to *C. formosanus* workers, causing 100% mortality after 24 h at the higher concentrations used; median concentrations caused lower mortality, but left survivors immobilized (Cornelius *et al.*, 1995). Even at lower concentrations, mortality of extract-exposed termites was three times that of controls after 48 h. Again, treated sand barriers were used, and showed that the repellent components of *O. glaber* extract (identified as two isomers of the monoterpene dolichodial) would significantly reduce the extent of tunnelling as long as 30 days after treatment.

In related work, extracellular metabolites (siderophores) of the brown-rot wood decay fungus *Gloeophyllum trabeum*

(Persoon: Fries) Murrill (Polyporaceae) were found to inhibit feeding by *C. formosanus* (Grace *et al.*, 1992). Siderophore-treated filter paper disks showed negligible feeding, whereas untreated disks were almost completely consumed over a 3-day test period.

Results of these and earlier studies (e.g. Amburgey & Beal, 1977) suggest that natural products, such as ant semiochemicals and fungal metabolites, or their synthetic analogues, might one day be of value in termite control programmes as repellents or insecticides in wood treatments or soil applications. However, the development of more stable formulations, such as microencapsulation, would be necessary to ensure their long-term, residual action.

Future research should establish what role biological methods, particularly the use of pathogens, effectively will play in overall termite management strategies. Biological control may yet come to supplement, but is unlikely any time soon to supplant, other established subterranean termite control technologies.

#### Acknowledgements

We thank S.Y. Higa (US Department of the Navy) and N.-Y. Su (University of Florida) for their critical reviews of an earlier draft of the manuscript, and Y. Liu for translation of Chinese material. Efforts of the second author (J.K. Grace) were partially supported by McIntire-Stennis funds and by USDA-ARS Specific Cooperative Agreements 58-6615-4-037 and 58-6615-9-018. This is Journal Series No. 4454 of the College of Tropical Agriculture and Human Resources, University of Hawaii.

#### References

- Al Fazairy, A.A. & Hassan, F.A. (1988) Infection of termites by *Spodoptera littoralis* nuclear polyhedrosis virus. *Insect Science and its Application* 9, 37-39.
- Al Fazairy, A.A. & Hassan, F.A. (1993) Histopathology of termite *Kaloterme flavicollis* Fabr. infected with a nuclear polyhedrosis virus. *Insect Science and its Application* 14, 127-134.
- Alton, R.A. (1947) A fungus parasitic on *Coptotermes curvignathus*, Holmgr. *Nature* 160, 120.
- Amburgey, T.L. & Beal, R.H. (1977) White rot inhibits termite attack. *Sociobiology* 3, 35-38.
- Arnold, G. (1915-1924) A monograph of the Formicidae of South Africa. *Annals of the South African Museum* 14, 1-766.
- Askew, R.R. (1971) *Parasitic insects*. 316 pp. New York, American Elsevier Publishing Company.
- Bao, L.-L. & Yendol, W.G. (1971) Infection of the eastern subterranean termite, *Reticulitermes flavipes* (Kollar) [sic] with the fungus *Beauveria bassiana* (Balsamo) Vuill. *Entomophaga* 16, 343-352.
- Beal, R.H. & Kais, A.G. (1962) Apparent infection of subterranean termites by *Aspergillus flavus* Link. *Journal of Insect Pathology* 4, 488-489.
- Becker, G. & Kerner-Gang, W. (1964) Schädigung und Förderung von Termiten durch Schimmelpilze. *Zeitschrift für Angewandte Entomologie* 53, 429-448.
- Blackwell, M. (1980) New records of termite-infesting fungi. *Journal of Invertebrate Pathology* 33, 101-104.
- Blackwell, M. & Gilbertson, R.L. (1981) *Cordycepsoides octosporus*, a termite suspected pathogen from Jalisco, Mexico. *Mycologia* 73, 358-362.
- Blackwell, M. & Gilbertson, R.L. (1984) New information on *Cordycepsoides bisporus* and *Cordycepsoides octosporus*. *Mycologia* 76, 763-765.
- Blackwell, M. & Kimbrough, J.W. (1976) Ultrastructure of the termite-associated fungus *Laboulbeniopsis termitarius*. *Mycologia* 68, 541-550.
- Blackwell, M. & Kimbrough, J.W. (1978) *Hormiscioideus filamentosus* gen. et sp. nov., a termite-infesting fungus from Brazil. *Mycologia* 70, 1274-1280.
- Blackwell, M. & Rossi, W. (1986) Biogeography of fungal ectoparasites of termites. *Mycotaxon* 25, 581-601.
- Bodot, P. (1961) La destruction des termitières de *Bellicositermes natalensis* Hav., par une Fourmi, *Dorylus* (*Typhlopone*) *dentifrons* Wasmann. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* 253, 3053-3054.
- Bridwell, J.C. (1920) Some notes on Hawaiian and other Bethyliidae [sic] (Hymenoptera) with the description of a new genus and species. 2nd paper. *Proceedings of the Hawaiian Entomological Society* 4, 291-305.
- Buchli, H. (1960) L'effet du champignon parasite *Antennopsis gallica* sur les jeunes colonies de Termites. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* 250, 1320-1321.
- Burges, H.D. (1981) Strategy for the microbial control of pests in 1980 and beyond. pp. 797-836 in Burges, H.D. (Ed.) *Microbial control of pests and plant diseases 1970-1980*. London, Academic Press.
- Colla, S. (1928) Su alcuni funghi parassiti delle termiti. *Bollettino del Laboratorio di Zoologia Generale e Agraria del R. Istituto Superiore Agrario di Portici* 22, 39-48.
- Collins, N.M. (1981) Populations, age structure and survivorship of colonies of *Macrotermes bellicosus* (Isoptera: Macrotermitinae). *Journal of Animal Ecology* 50, 293-311.
- Cornelius, M.L. & Grace, J.K. (1994) Semiochemicals extracted from a dolichoderine ant affects [sic] the feeding and tunneling behavior of the Formosan subterranean termite (Isoptera: Rhinotermitidae). *Journal of Economic Entomology* 87, 705-708.
- Cornelius, M.L. & Grace, J.K. (1996) Effect of two ant species (Hymenoptera: Formicidae) on the foraging and survival of the Formosan subterranean termite (Isoptera: Rhinotermitidae). *Environmental Entomology* 25, 85-89.
- Cornelius, M.L. & Grace, J.K. (1997) Effect of termite soldiers on the foraging behavior of *Coptotermes formosanus* (Isoptera: Rhinotermitidae) in the presence of predatory ants. *Sociobiology* 29, 247-253.
- Cornelius, M.L., Grace, J.K., Ford, P.W. & Davidson, B.S. (1995) Toxicity and repellency of semiochemicals extracted from a dolichoderine ant (Hymenoptera: Formicidae) to the Formosan subterranean termite (Isoptera: Rhinotermitidae). *Environmental Entomology* 24, 1263-1269.
- Danthanarayana, W. & Vitarana, S.I. (1987) Control of the live-wood tea termite *Glyptotermes dilatatus* using *Heterorhabditis* sp. (Nemat.). *Agriculture, Ecosystems and Environment* 19, 333-342.
- DeBach, P. & Rosen, D. (1991) *Biological control by natural enemies*. 2nd edn. 440 pp. Cambridge, Cambridge University Press.
- Delate, K.M., Grace, J.K. & Tome, C.H.M. (1995) Potential use of pathogenic fungi in baits to control the Formosan subterranean termite (Isopt., Rhinotermitidae). *Journal of Applied Entomology* 119, 429-433.
- Deligne, J., Quennedey, A. & Blum, M.S. (1981) The enemies



- and defense mechanisms of termites. pp. 1-76 in Hermann, H.R. (Ed.) *Social insects*, vol. 2. New York, Academic Press.
- Desportes, I. (1963) Cycle évolutif d'une nouvelle Grégarine parasite de Termites: *Diplocystis zootermopsidis* sp.n. (Eugregarina, Diplocystidae). *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* 257, 4013-4015.
- Disney, R.H.L. (1986) Two remarkable new species of scuttle-fly (Diptera: Phoridae) that parasitize termites (Isoptera) in Sulawesi. *Systematic Entomology* 11, 413-422.
- Disney, R.H.L. (1988) Biology and taxonomy of old world *Puliciphora* (Diptera: Phoridae) with descriptions of nine new species. *Oriental Insects* 22, 267-286.
- Disney, R.H.L. & Kistner, D.H. (1989) Six new species of oriental Phoridae from termite nests (Diptera; Isoptera; Termitidae and Rhinotermitidae). *Sociobiology* 13, 325-348.
- Disney, R.H.L. & Kistner, D.H. (1990) Revision of the termite-parasitizing genus *Dicranopteron* (Diptera: Phoridae). *Zoological Journal of the Linnean Society* 98, 111-122.
- Edwards, R. & Mill, A.E. (1986) *Termites in buildings: their biology and control*. 261 pp. East Grinstead, United Kingdom, Rentokil Ltd.
- Epsky, N.D. & Capinera, J.L. (1988) Efficacy of the entomogenous nematode *Steinernema feltiae* against a subterranean termite, *Reticulitermes tibialis* (Isoptera: Rhinotermitidae). *Journal of Economic Entomology* 81, 1313-1317.
- Federici, B.A. (1990) Bright horizons for invertebrate pathology. pp. v-ix in *Proceedings and Abstracts. Vth International Colloquium on Invertebrate Pathology and Microbial Control, Adelaide, Australia, 20-24 August 1990*. Adelaide, Society for Invertebrate Pathology.
- Fernandes, P.M. (1991) Controle microbiano de *Cornitermes cumulans* (Kollar, 1832) utilizando *Beauveria bassiana* (Bals.) Vuill. e *Metarhizium anisopliae* (Metsch.) Sorok. PhD dissertation, ESALQ-Universidade de Sao Paulo, Piracicaba, Brazil.
- Ferron, P. (1978) Biological control of insect pests by entomogenous fungi. *Annual Review of Entomology* 23, 409-442.
- Fujii, J.K. (1975) Effects of an entomogenous nematode, *Neoaplectana carpocapsae* Weiser, on the Formosan subterranean termite, *Coptotermes formosanus* Shiraki, with ecological and biological studies on *C. formosanus*. 163 pp. PhD dissertation, University of Hawaii, Honolulu, Hawaii, USA.
- Fuxa, J.R. (1990) New directions for insect control with baculoviruses. pp. 97-113 in Baker, R.R. & Dunn, P.E. (Eds) *New directions in biological control: alternatives for suppressing agricultural pests and diseases*. New York, Alan R. Liss, Inc.
- Gao, D.-R. (1987) Use of attractants in bait toxicants for the control of *Coptotermes formosanus* Shiraki in China. pp. 53-57 in Tamashiro, M. & Su, N.-Y. (Eds) *Biology and control of the Formosan subterranean termite*. Honolulu, University of Hawaii College of Tropical Agriculture and Human Resources Research Extension Series 083.
- Gaugler, R. (1988) Ecological considerations in the biological control of soil-inhabiting insects with entomopathogenic nematodes. *Agriculture, Ecosystems and Environment* 24, 351-360.
- Georgis, R., Poinar, G.O., Jr. & Wilson, A.P. (1982) Susceptibility of damp-wood termites and soil and wood-dwelling termites to the entomogenous nematode *Neoaplectana carpocapsae*. *IRCS Medical Science* 10, 563.
- Gibbs, A.J., Gay, F.J. & Wetherly, A.H. (1970) A possible paralysis virus of termites. *Virology* 40, 1063-1065.
- Gitonga, W. (1996) *Metarhizium anisopliae* (Metschnikoff) Sorokin and *Beauveria bassiana* (Balsamo) Vuillemin as potential biological control agents of *Macrotermes michaelseni* (Sjöstedt) (Isoptera: Termitidae) in Kenya. 144 pp. PhD dissertation, Royal Veterinary and Agricultural University, Copenhagen, Denmark.
- Gouger, R.J. & Kimbrough, J.W. (1969) *Antennopsis gallica* Heim and Buchli (Hyphomycetes: Gloeochaustoriales), an entomogenous fungus on subterranean termites in Florida. *Journal of Invertebrate Pathology* 13, 223-228.
- Grace, J.K. (1991) Termite-fungal associations and manipulations for termite control. p. 29 in *Program and Abstracts. 24th annual meeting, Society for Invertebrate Pathology, Northern Arizona University, August 4-9, 1991*. Flagstaff, Arizona, Society for Invertebrate Pathology.
- Grace, J.K. (1993) Microbial termite control: effects of entomogenous fungi on the Formosan subterranean termite (Isoptera: Rhinotermitidae). p. 474 in Wilder, K.B. & Robinson, W.H. (Eds) *Proceedings of the 1st International Conference on Insect Pests in the Urban Environment, St John's College, Cambridge, 30 June-3 July 1993*. Exeter, Organizing Committee of the International Conference on Insect Pests in the Urban Environment.
- Grace, J.K. (1994) Protocol for testing effects of microbial pest control agents on nontarget subterranean termites (Isoptera: Rhinotermitidae). *Journal of Economic Entomology* 87, 269-274.
- Grace, J.K. (1995) Microbial termite control. pp. 166-167 in *Hawaii agriculture: positioning for growth. Proceedings of a conference organized by the College of Tropical Agriculture & Human Resources, University of Hawaii at Manoa, 5-6 April 1995*. Honolulu, CTAHR.
- Grace, J.K. (1997) Biological control strategies for suppression of termites. *Journal of Agricultural Entomology* 14, 281-289.
- Grace, J.K. & Ewart, D.M. (1996) Recombinant cells of *Pseudomonas fluorescens*: a highly palatable encapsulation for delivery of genetically engineered toxins to subterranean termites (Isoptera: Rhinotermitidae). *Letters in Applied Microbiology* 23, 183-186.
- Grace, J.K. & Zoberi, M.H. (1992) Experimental evidence for transmission of *Beauveria bassiana* by *Reticulitermes flavipes* workers (Isoptera: Rhinotermitidae). *Sociobiology* 20, 23-28.
- Grace, J.K., Goodell, B.S., Jones, W.E., Chandhoke, V. & Jellison, J. (1992) Evidence for inhibition of termite (Isoptera: Rhinotermitidae) feeding by extracellular metabolites of a wood decay fungus. *Proceedings of the Hawaiian Entomological Society* 31, 249-252.
- Grace, J.K., Tome, C.H.M., Shelton, T.G., Oshiro, R.J. & Yates, J.R. III (1996) Baiting studies and considerations with *Coptotermes formosanus* (Isoptera: Rhinotermitidae) in Hawaii. *Sociobiology* 28, 511-520.
- Gross, P. (1991) Influence of target pest feeding niche on success rates in classical biological control. *Environmental Entomology* 20, 1217-1227.
- Hänel, H. (1981) A bioassay for measuring the virulence of the insect pathogenic fungus *Metarhizium anisopliae* (Metsch.) Sorok. (fungi imperfecti) [sic] against the termite *Nasutitermes exitiosus* (Hill) (Isoptera, Termitidae). *Zeitschrift für Angewandte Entomologie* 92, 9-18.
- Hänel, H. (1982a) The life cycle of the insect pathogenic fungus *Metarhizium anisopliae* in the termite *Nasutitermes exitiosus*. *Mycopathologia* 80, 137-145.

- Hänel, H. (1982b) Selection of a fungus species, suitable for the biological control of the termite *Nasutitermes exitiosus* (Hill). *Zeitschrift für Angewandte Entomologie* 94, 237-245.
- Hänel, H. & Watson, J.A.L. (1983) Preliminary field tests on the use of *Metarhizium anisopliae* for the control of *Nasutitermes exitiosus* (Hill) (Isoptera: Termitidae). *Bulletin of Entomological Research* 73, 305-313.
- HDOA (1994) Report to the Eighteenth Legislature, 1995 regular session, in response to Section 20.1, Act 252, Session Laws of Hawaii, 1994 regular session, requesting the Department of Agriculture to evaluate methods for controlling the Formosan termite through biological control agents. 24 pp. Unpublished report. Honolulu, Hawaii Department of Agriculture.
- Henry, J.E. (1990) Control of insects by Protozoa. pp. 161-176 in Baker, R.R. & Dunn, P.E. (Eds) *New directions in biological control: alternatives for suppressing agricultural pests and diseases*. New York, Alan R. Liss, Inc.
- Hölldobler, B. & Wilson, E.O. (1990) *The ants*. 732 pp. Cambridge, Massachusetts, Belknap Press of Harvard University Press.
- Howarth, F.G. (1991) Environmental impacts of classical biological control. *Annual Review of Entomology* 36, 485-509.
- Huffaker, C.B. & Messenger, P.S. (Eds) (1976) *Theory and practice of biological control*. 788 pp. New York, Academic Press.
- Jafri, R.H., Ahmad, M. & Idrees, K. (1976) Microsporidian infection in the workers of termite, *Microcerotermes championi*. *Pakistan Journal of Zoology* 8, 234-236.
- Johnson, J.B. & Hagen, K.S. (1981) A neuropterous larva uses an allomone to attack termites. *Nature* 289, 506-507.
- Jones, W.E., Grace, J.K. & Tamashiro, M. (1996) Virulence of seven isolates of *Beauveria bassiana* and *Metarhizium anisopliae* to *Coptotermes formosanus* (Isoptera: Rhinotermitidae). *Environmental Entomology* 25, 481-487.
- Kaya, H.K. (1990) Entomopathogenic nematodes in biological control of insects. pp. 189-198 in Baker, R.R. & Dunn, P.E. (Eds) *New directions in biological control: alternatives for suppressing agricultural pests and diseases*. New York, Alan R. Liss, Inc.
- Kaya, H.K. & Gaugler, R. (1993) Entomopathogenic nematodes. *Annual Review of Entomology* 38, 181-206.
- Kaya, H.K., Bedding, R.A. & Akhurst, R.J. (1993) An overview of insect-parasitic and entomopathogenic nematodes. pp. 1-10 in Bedding, R.A., Akhurst, R. & Kaya, H. (Eds) *Nematodes and the biological control of insect pests*. East Melbourne, Australia, CSIRO.
- Kevoorkian, A.G. (1937) Studies in the Entomophthoraceae. I. Observations on the genus *Conidiobolus*. *Journal of Agriculture of the University of Puerto Rico* 21, 191-200.
- Khan, K.I., Fazal, Q. & Jafri, R.H. (1977a) Pathogenicity of locally discovered *Bacillus thuringiensis* strain to the termites: *Heterotermes indicola* (Wassman) [sic] and *Microcerotermes championi* (Snyder). *Pakistan Journal of Scientific Research* 29, 12-13.
- Khan, K.I., Fazal, Q., Jafri, R.H. & Ahmad, M. (1977b) Susceptibility of various species of termites to a pathogen, *Serratia marcescens*. *Pakistan Journal of Scientific Research* 29, 46-47.
- Khan, K.I., Fazal, Q. & Jafri, R.H. (1978) Development of *Bacillus thuringiensis* in a termite, *Heterotermes indicola* (Wassman) [sic]. *Pakistan Journal of Science* 30, 117-119.
- Khan, K.I., Jafri, R.H. & Ahmad, M. (1985) The pathogenicity and development of *Bacillus thuringiensis* in termites. *Pakistan Journal of Zoology* 17, 201-209.
- Khan, K.I., Jafri, R.H., Ahmad, M. & Khan, K.M.S. (1992) The pathogenicity of *Pseudomonas aeruginosa* against termites. *Pakistan Journal of Zoology* 24, 243-245.
- Khan, S.R. & Kimbrough, J.W. (1974a) Morphology and development of *Termitaria snyderi*. *Mycologia* 66, 446-462.
- Khan, S.R. & Kimbrough, J.W. (1974b) Taxonomic position of *Termitaria* and *Mattirolella* (entomogenous Deuteromycetes). *American Journal of Botany* 61, 395-399.
- Kimbrough, J.W. & Gouger, R.J. (1970) Structure and development of the fungus *Laboulbeniopsis termitarius*. *Journal of Invertebrate Pathology* 16, 205-213.
- Kimbrough, J.W., Morales, M.I. & Gouger, R.J. (1972) A new species of *Dimeromyces* (Laboulbeniales) on subterranean termites. *Mycologia* 64, 388-393.
- Klein, M.G. (1988) Pest management of soil-inhabiting insects with microorganisms. *Agriculture, Ecosystems and Environment* 24, 337-349.
- Klein, M.G. (1990) Efficacy against soil-inhabiting insect pests. pp. 195-214 in Gaugler, R. & Kaya, H.K. (Eds) *Entomopathogenic nematodes in biological control*. Boca Raton, Florida, CRC Press.
- Ko, W.H., Fujii, J.K. & Kanegawa, K.M. (1982) The nature of soil pernicious to *Coptotermes formosanus*. *Journal of Invertebrate Pathology* 39, 38-40.
- Kramm, K.R. & West, D.F. (1982) Termite pathogens: effects of ingested *Metarhizium*, *Beauveria*, and *Gliocladium* conidia on worker termites (*Reticulitermes* sp.). *Journal of Invertebrate Pathology* 40, 7-11.
- Kramm, K.R., West, D.F. & Rockenbach, P.G. (1982) Termite pathogens: transfer of the entomopathogen *Metarhizium anisopliae* between *Reticulitermes* sp. termites. *Journal of Invertebrate Pathology* 40, 1-6.
- Kühne, H. & Becker, G. (1974) Zur Biologie und Ökologie von *Scleroderma domesticum* Latreille (Bethyilidae, Hymenoptera), einem Parasiten holzzerstörender Insektenlarven. *Zeitschrift für Angewandte Entomologie* 76, 278-303.
- Lai, P.-Y. (1977) Biology and ecology of the Formosan subterranean termite, *Coptotermes formosanus*, and its susceptibility to the entomogenous fungi, *Beauveria bassiana* and *Metarhizium* [sic] *anisopliae*. 140 pp. PhD dissertation, University of Hawaii, Honolulu, Hawaii, USA.
- Lai, P.Y., Tamashiro, M. & Fujii, J.K. (1982) Pathogenicity of six strains of entomogenous fungi to *Coptotermes formosanus*. *Journal of Invertebrate Pathology* 39, 1-5.
- Lai, P.Y., Tamashiro, M., Yates, J.R., Su, N.Y., Fujii, J.K. & Ebesu, R.H. (1983) Living plants in Hawaii attacked by *Coptotermes formosanus*. *Proceedings of the Hawaiian Entomological Society* 24, 283-286.
- Laumond, C., Mauléon, H. & Kermarrec, A. (1979) Données nouvelles sur le spectre d'hôtes et le parasitisme du nématode entomophage *Neoplectana carpocapsae*. *Entomophaga* 24, 13-27.
- Lee, K.E. & Wood, T.G. (1971) *Termites and soils*. 251 pp. London, Academic Press.
- Leong, K.L.H. (1966) Infection of the Formosan subterranean termite, *Coptotermes formosanus* Shiraki [sic] by the fungus, *Metarhizium* [sic] *anisopliae* (Metch.) Sorokin. 53 pp. Master's thesis, University of Hawaii, Honolulu, Hawaii, USA.
- Lepage, M.G. & Darlington, J.P.E.C. (1984) Observations on the ant *Carebara vidua* F. Smith preying on termites in Kenya. *Journal of Natural History* 18, 293-302.
- Levieux, J. (1966) Note préliminaire sur les colonnes de chasse

- de *Megaponera foetens* F. (Hyménoptère Formicidae). *Insectes Sociaux* 13, 117-126.
- Lin, S.-Q. (1987) Present status of *Coptotermes formosanus* and its control in China. pp. 31-37 in Tamashiro, M. & Su, N.-Y. (Eds) *Biology and control of the Formosan subterranean termite*. Honolulu, University of Hawaii College of Tropical Agriculture and Human Resources Research Extension Series 083.
- Logan, J.W.M., Cowie, R.H. & Wood, T.G. (1990) Termite (Isoptera) control in agriculture and forestry by non-chemical methods: a review. *Bulletin of Entomological Research* 80, 309-330.
- Longhurst, C. & Howse, P.E. (1979) Foraging, recruitment and emigration in *Megaponera foetens* (Fab.) (Hymenoptera: Formicidae) from the Nigerian Guinea Savanna. *Insectes Sociaux* 26, 204-215.
- Longhurst, C., Johnson, R.A. & Wood, T.G. (1978) Predation by *Megaponera foetens* (Fabr.) (Hymenoptera: Formicidae) on termites in the Nigerian southern Guinea Savanna. *Oecologia* 32, 101-107.
- Longhurst, C., Baker, R. & Howse, P.E. (1979a) Termite predation by *Megaponera foetens* (Fab.) (Hymenoptera: Formicidae): coordination of raids by glandular secretions. *Journal of Chemical Ecology* 5, 703-719.
- Longhurst, C., Johnson, R.A. & Wood, T.G. (1979b) Foraging, recruitment and predation by *Decamorium uelense* (Sanstch) (Formicidae: Myrmicinae) on termites in southern Guinea Savanna, Nigeria. *Oecologia* 38, 83-91.
- Lund, A.E. (1971) Microbial control of termites. pp. 385-386 in Burges, H.D. & Hussey, N.W. (Eds) *Microbial control of insects and mites*. London, Academic Press.
- Lund, A.E. & Engelhardt, N.T. (1962) Subterranean termites and *Absidia coerulea* Bainier (Mucorales). *Journal of Insect Pathology* 4, 131-132.
- Martin, P.A.W. (1994) An iconoclastic view of *Bacillus thuringiensis* ecology. *American Entomologist* 40, 85-90.
- Maschwitz, U. & Mühlberg, M. (1975) Zur Jagdstrategie einiger orientalischer *Leptogenys*-Arten (Formicidae: Ponerinae). *Oecologia* 20, 65-83.
- Maschwitz, U. & Schönegge, P. (1983) Forage communication, nest moving recruitment, and prey specialization in the oriental ponerine *Leptogenys chinensis*. *Oecologia* 57, 175-182.
- Mauldin, J.K. (1986) Economic importance and control of termites in the United States. pp. 130-143 in Vinson, S.B. (Ed.) *Economic impact and control of social insects*. New York, Praeger Publishers.
- Mauldin, J.K. & Beal, R.H. (1989) Entomogenous nematodes for control of subterranean termites, *Reticulitermes* spp. (Isoptera: Rhinotermitidae). *Journal of Economic Entomology* 82, 1638-1642.
- McCoy, C.W. (1990) Entomogenous fungi as microbial pesticides. pp. 139-159 in Baker, R.R. & Dunn, P.E. (Eds) *New directions in biological control: alternatives for suppressing agricultural pests and diseases*. New York, Alan R. Liss, Inc.
- Merrill, J.H. & Ford, A.L. (1916) Life history and habits of two new nematodes parasitic on insects. *Journal of Agricultural Research* 6, 115-127.
- Milner, R.J. (1992) Selection and characterization of strains of *Metarhizium anisopliae* for control of soil insects in Australia. pp. 200-207 in Lomer, C.J. & Prior, C. (Eds) *Biological control of locusts and grasshoppers*. Wallingford, CAB International.
- Milner, R.J., Staples, J.A. & Lenz, M. (1996) Options for termite management using the insect pathogenic fungus *Metarhizium anisopliae*. *International Research Group on Wood Preservation*. Document No. IRG/WP 9610142, pp. 1-5.
- Mori, H. (1987) The Formosan subterranean termite in Japan: its distribution, damage, and current and potential control measures. pp. 23-26 in Tamashiro, M. & Su, N.-Y. (Eds) *Biology and control of the Formosan subterranean termite*. Honolulu, University of Hawaii College of Tropical Agriculture and Human Resources Research Extension Series 083.
- Nickle, W.R. & Welch, H.E. (1984) History, development, and importance of insect nematology. pp. 627-653 in Nickle, W.R. (Ed.) *Plant and insect nematodes*. New York, Marcel Dekker, Inc.
- Nutting, W.L. (1969) Flight and colony foundation. pp. 233-282 in Krishna, K. & Weesner, F.M. (Eds) *Biology of termites*, vol. 1. New York, Academic Press.
- Ochiel, G.R.S. (1995) Biology and biocontrol potential of *Cordycepioideus bisporus* Stifter and *Paecilomyces fumosoroseus* (Wize) Brown and Smith on the higher termite *Macrotermes subhyalinus* (Rambur) in Kenya. 210 pp. PhD dissertation, Royal Veterinary and Agricultural University, Copenhagen, Denmark.
- Ochiel, G.R.S., Evans, H.C. & Ellenberg, J. (1997) *Cordycepioideus*, a pathogen of termites in Kenya. *Mycologist* 11, 7-9.
- Ohlgu, C.E. & Wood, T.G. (1976) A method for measuring rate of grass-harvesting by *Trinervitermes geminatus* (Wasmann) (Isoptera, Nasutitermitinae) and observation on its foraging behaviour in southern Guinea Savanna, Nigeria. *Journal of Applied Ecology* 13, 705-713.
- Pearce, M.J. (1997) *Termites: biology and pest management*. 172 pp. Wallingford, CAB International.
- Pemberton, C.E. (1928a) Thysanuran predatory on eggs and immature forms of termites in Borneo. *Proceedings of the Hawaiian Entomological Society* 7, 147.
- Pemberton, C.E. (1928b) Nematodes associated with termites in Hawaii, Borneo and Celebes. *Proceedings of the Hawaiian Entomological Society* 7, 148-150.
- Phillipsen, W.J. & Coppel, H.C. (1977) *Acotyledon formosani* sp. n. associated with the Formosan subterranean termite, *Coptotermes formosanus* Shiraki (Acarina: Acaridae-Isoptera: Rhinotermitidae). *Journal of the Kansas Entomological Society* 50, 399-409.
- Pimentel, D., Glenister, C., Fast, S. & Gallahan, D. (1984) Environmental risks of biological pest controls. *Oikos* 42, 283-290.
- Poptiel, I. & Hominick, W.H. (1992) Nematodes as biological control agents: Part II. *Advances in Parasitology* 31, 381-433.
- Preston, A.F., Erbsch, F.H., Kramm, K.R. & Lund, A.E. (1982) Developments in the use of biological control for wood preservation. *Proceedings of the American Wood-Preservers' Association* 78, 53-61.
- Rath, A.C. & Tidbury, C.A. (1996) Susceptibility of *Coptotermes acinaciformis* (Isoptera: Rhinotermitidae) and *Nasutitermes exitiosus* (Isoptera: Termitidae) to two commercial isolates of *Metarhizium anisopliae*. *Sociobiology* 28, 67-72.
- Rosengaus, R.B. & Traniello, J.F.A. (1997) Pathobiology and disease transmission in dampwood termites [*Zootermopsis angusticollis* (Isoptera: Termopsidae)] infected with the fungus *Metarhizium anisopliae* (Deuteromycotina: Hypomycetes). *Sociobiology* 30, 185-195.
- Rossi, W. (1974) Una nuova specie di *Laboulbenia* (Ascomycetes, Laboulbeniales) parassita di termiti. *Doriana* 5, 1-5.
- Sajap, A.S. & Kaur, K. (1990) Histopathology of *Metarhizium*

- anisopliae*, an entomopathogenic fungus, infection in the termite, *Coptotermes curvignathus*. *Pertanika* 13, 331-334.
- Sajap, A.S., Atim, A.B., Husin, H. & Wahab, Y.A. (1997) Isolation of *Conidiobolus coronatus* (Zygomycetes: Entomophthorales) from soil and its effect on *Coptotermes curvignathus* (Isoptera: Rhinotermitidae). *Sociobiology* 30, 257-262.
- Sannasi, A. (1968) Apparent infection of queens and drones of the mound-building termite *Odontotermes obesus* by *Aspergillus flavus*. *Journal of Invertebrate Pathology* 10, 434-435.
- Schmitz, H. (1938) *Misotermes exenterans* n.g. n.sp., eine parasitische Fliege aus der Familie der Phoridae, welche die Entstehung myliagener Soldaten bei *Macrotermes gilvius* Hag. in Java verursacht. *Treubia* 16, 369-397.
- Secord, D. & Kareiva, P. (1996) Perils and pitfalls in the host specificity paradigm. *BioScience* 46, 448-453.
- Sheppe, W. (1970) Invertebrate predation on termites of the African savanna. *Insectes Sociaux* 17, 205-218.
- Smith, K.M. (1967) *Insect virology*. 256 pp. New York, Academic Press.
- Smythe, R.V. & Coppel, H.C. (1965) The susceptibility of *Reticulitermes flavipes* (Kollar) and other termite species to an experimental preparation of *Bacillus thuringiensis* Berliner. *Journal of Invertebrate Pathology* 7, 423-426.
- Smythe, R.V. & Coppel, H.C. (1966) Pathogenicity of externally occurring fungi to *Reticulitermes flavipes*. *Journal of Invertebrate Pathology* 8, 266-267.
- Staples, J. & Milner, R. (1996) *Metarhizium anisopliae* as a mycotermiticide-laboratory behavioural bioassay. p. 493 in Wildey, K.B. (Ed.) *Proceedings of the 2nd International Conference on Insect Pests in the Urban Environment, Edinburgh, 7-10 July 1996*. Exeter, Organizing Committee of the International Conference on Insect Pests in the Urban Environment.
- Steinhaus, E.A. & Marsh, G.A. (1962) Report of diagnoses of diseased insects 1951-1961. *Hilgardia* 33, 349-490.
- Stiffler, C.B. (1941) A new genus of Hypocreales. *Mycologia* 33, 82-86.
- Su, N.-Y. (1993) Managing subterranean termite populations. pp. 45-50 in Wildey, K.B. & Robinson, W.H. (Eds) *Proceedings of the 1st International Conference on Insect Pests in the Urban Environment, Cambridge, 30 June-3 July 1993*. Exeter, Organizing Committee of the International Conference on Insect Pests in the Urban Environment.
- Su, N.-Y. (1994) Field evaluation of a hexaflumuron bait for population suppression of subterranean termites (Isoptera: Rhinotermitidae). *Journal of Economic Entomology* 87, 389-397.
- Su, N.-Y. & Scheffrahn, R.H. (1990) Economically important termites in the United States and their control. *Sociobiology* 17, 77-94.
- Su, N.-Y. & Scheffrahn, R.H. (1993) Laboratory evaluation of two chitin synthesis inhibitors, hexaflumuron and diflubenzuron, as bait toxicants against Formosan and eastern subterranean termites (Isoptera: Rhinotermitidae). *Journal of Economic Entomology* 86, 1453-1457.
- Su, N.-Y. & Scheffrahn, R.H. (1998) A review of subterranean termite control practices and prospects for integrated pest management programmes. *Integrated Pest Management Reviews* 3, 1-13.
- Su, N.-Y. & Tamashiro, M. (1987) An overview of the Formosan subterranean termite (Isoptera: Rhinotermitidae) in the world. pp. 3-15 in Tamashiro, M. & Su, N.-Y. (Eds) *Biology and control of the Formosan subterranean termite*. Honolulu, University of Hawaii College of Tropical Agriculture and Human Resources Research Extension Series 083.
- Su, N.-Y., Thoms, E.M., Ban, P.M. & Scheffrahn, R.H. (1995) Monitoring/baiting station to detect and eliminate foraging populations of subterranean termites (Isoptera: Rhinotermitidae) near structures. *Journal of Economic Entomology* 88, 932-936.
- Suzuki, K. (1991) Laboratory trials of biological control agents against subterranean termites. *International Research Group on Wood Preservation*. Document No. IRG/WP/1475, pp. 1-11.
- Suzuki, K. (1996) Biological control of termites by pathogenic fungi. pp. 146-156 in *Forestry and Forest Products Research: Proceedings of the Third Conference, Kepong, Malaysia, 3-4 October 1995, volume 2*. Kepong, Forest Research Institute of Malaysia.
- Tamashiro, M., Yates, J.R. & Ebesu, R.H. (1987) The Formosan subterranean termite in Hawaii: problems and control. pp. 15-22 in Tamashiro, M. & Su, N.-Y. (Eds) *Biology and control of the Formosan subterranean termite*. Honolulu, University of Hawaii College of Tropical Agriculture and Human Resources Research Extension Series 083.
- Tate, P. (1927) On *Ectomyces calotermiti* n.g., n.sp., an Ascomycete parasitic on *Calotermes samoanus* Holmgren (Isoptera, Protermitidae). *Parasitology* 19, 54-60.
- Tate, P. (1928) Notes on the genera *Ectomyces* and *Termitaria*, fungi parasitic on termites. *Parasitology* 20, 77-78.
- Thaxter, R. (1895) Notes on Laboulbeniaceae, with descriptions of new species. *Proceedings of the American Academy of Arts and Sciences* 30, 467-481.
- Thaxter, R. (1920) Second note on certain peculiar fungus-parasites of living insects. *Botanical Gazette* 69, 1-27.
- Toumanoff, C. (1965) Action de divers champignons entomophages sur *Reticulitermes santonensis*: aperçu général. *Annales de Parasitologie Humaine et Comparée* 40, 611-624.
- Toumanoff, C. & Rombaut, J. (1965) Action de certains champignons entomophages, cultivés sur les milieux appropriés attractifs, sur le Termeite de Saintonge *Reticulitermes santonensis* (de Feytaud). *Annales de Parasitologie Humaine et Comparée* 40, 605-609.
- Traniello, J.F.A. (1981) Enemy deterrence in the recruitment strategy of a termite: soldier-organized foraging in *Nasutitermes costalis*. *Proceedings of the National Academy of Sciences of the United States of America* 78, 1976-1979.
- Trudeau, D. (1989) Selection of entomophilic nematodes for control of the eastern subterranean termite, *Reticulitermes flavipes* (Kollar) (Isoptera: Rhinotermitidae). 93 pp. Master's thesis, University of Toronto, Toronto, Ontario, Canada.
- USDA (1999) Agricultural Research Service National Formosan Termite Program; <http://www.ars.usda.gov/is/fullstop/faqenviro.htm>; 3 pp. [Accessed 15 September 1999].
- Veeresh, G.K. & Gubbalah. (1984) A report on the 'crazy ant' (*Anoplolepis longipes* Jerdon) menace in Karnataka. *Journal of Soil Biology and Ecology* 4, 65-73.
- Wahlman, M. & Davidson, B.S. (1993) New destruxins from the entomopathogenic fungus *Metarhizium anisopliae*. *Journal of Natural Products (Lloydia)* 56, 643-647.
- Weber, N.A. (1964) Termite prey of some African ants. *Entomological News* 75, 197-204.
- Wells, J.D. & Henderson, G. (1993) Fire ant predation on native and introduced subterranean termites in the laboratory:

- effect of high soldier number in *Coptotermes formosanus*. *Ecological Entomology* 18, 270-274.
- Wells, J.D., Fuxa, J.R. & Henderson, G. (1995) Virulence of four fungal pathogens to *Coptotermes formosanus* (Isoptera: Rhinotermitidae). *Journal of Entomological Science* 30, 208-215.
- Wheeler, W.M. (1936) Ecological relations of ponerine and other ants to termites. *Proceedings of the American Academy of Arts and Sciences* 71, 159-243.
- Williams, R.M.C. (1959) Colony development in *Cubitermes ugandensis* Fuller (Isoptera: Termitidae). *Insectes Sociaux* 6, 291-304.
- Wilson, N.L. & Oliver, A.D. (1969) Food habits of the imported fire ant in pasture and pine forest areas in southeastern Louisiana. *Journal of Economic Entomology* 62, 1268-1271.
- Wood, T.G. & Johnson, R.A. (1986) The biology, physiology, and ecology of termites. pp. 1-68 in Vinson, S.B. (Ed.) *Economic impact and control of social insects*. New York, Praeger Publishers.
- Wood, T.G. & Sands, W.A. (1978) The role of termites in ecosystems. pp. 245-292 in Brian, M.V. (Ed.) *Production ecology of ants and termites*. Cambridge, Cambridge University Press.
- Yates, J.R. III, Grace, J.K. & Tamashiro, M. (1997) The Formosan subterranean termite: a review of new management methods in Hawaii. pp. 59-68 in Chan, J. (Ed.) *Proceedings of 1997 FAOPMA Convention held in Hong Kong, April 7-9, 1997*. Hong Kong, Hong Kong Pest Control Association.
- Yendol, W.G. & Paschke, J.D. (1965) Pathology of an *Entomophthora* infection in the eastern subterranean termite *Reticulitermes flavipes* (Kollar). *Journal of Invertebrate Pathology* 7, 414-422.
- Yin, S.-C. (1994?) Inveterate enemy of subterranean termites *Dorylus orientalis* Westwood. p. 2333 in Xia, K. (Ed.) *Selected papers on Chinese termitology (1984-1993)*, vol. 3. Nanjing, Tian Ze Press.
- Yoshimura, T. & Takahashi, M. (1998) Termiticidal performance of an entomogenous fungus, *Beauveria brongniartii* (Saccardo) Petch [sic] in laboratory tests. *Japanese Journal of Environmental Entomology and Zoology* 9, 16-22.
- Yoshimura, T., Tsunoda, K., Takahashi, M. & Katsuda, Y. (1992) Pathogenicity of an entomogenous fungus, *Conidiobolus coronatus* Tyrrell and Macleod, to *Coptotermes formosanus* Shiraki. *Japanese Journal of Environmental Entomology and Zoology* 4, 11-16.
- Zimmerman, E.C. (1948) *Insects of Hawaii*, vol. 2. *Apterygota to Thysanoptera*. 475 pp. Honolulu, University of Hawaii Press.
- Zoberi, M.H. (1995) *Metarhizium anisopliae*, a fungal pathogen of *Reticulitermes flavipes* (Isoptera: Rhinotermitidae). *Mycologia* 87, 354-359.
- Zoberi, M.H. & Grace, J.K. (1990a) Isolation of the pathogen *Beauveria bassiana* from *Reticulitermes flavipes* (Isoptera: Rhinotermitidae). *Sociobiology* 16, 289-296.
- Zoberi, M.H. & Grace, J.K. (1990b) Fungi associated with the subterranean termite *Reticulitermes flavipes* in Ontario. *Mycologia* 82, 289-294.

(Accepted 1 November 1999)  
© CAB International, 2000