

College of Tropical Agriculture and Human Resources University of Hawai'i at Mānoa

Fruits and Nuts June 2011 F_N-21

Rollinia

Ken Love¹ and Robert E. Paull² ¹Hawaii Tropical Fruit Growers, ²CTAHR Department of Tropical Plant and Soil Sciences

Rollinia deliciosa Safford Rollinia mucosa (Jacq.) Baill Family: Annonaceae

R ollinia is a fast-growing tropical tree that can reach heights of 50 feet and thrives from 300 to 3000 feet elevation in Hawai'i, as long as it receives enough water. It will grow in a wide variety of deep soils and can gain as much as 10 feet in height per year. The tree will not tolerate cold or extended periods of drought. Under ideal conditions, the tree can fruit as

early as $2\frac{1}{2}$ years from the time of planting. The fruit is highly inconsistent in shape and size. It turns from green to yellow when ripe. The somewhat fragile skin contains milky-white, pyramidal shaped carpels usually containing black seeds that average half an inch in length.

Limited commercial planting means that most sales occur at small markets and farm stands. The fruit is rapidly gaining popularity as an income source for smallholder farmers in Brazil. In Hawai'i, the fruit can be found at many farmers' markets and health food stores and is rapidly gaining favor with chefs and larger groceries featuring locally grown produce.

Various references list *Rollinia mucosa* (Jacq.) Baill and *Rollinia deliciosa* Safford as two species, while many others list them as synonyms.

Other common names

biriba de Pernambuco, fruta da condessa, jaca de pobre, araticu, araticum, araticum pitaya (Brazil);



mulato (Colombia); rinon or rinon de monte (Venezuela); anona babosa, zambo (Mexico); anon cimarron, corosol, cachiman, cashina, candongo anon de monte, articum, condessa, graviolabrava; wild sweetsop, wild sugar apple (English)

Origin

Rollinia is thought to have originated in northern Brazil along the banks of the Amazon. It is also found in its natural state in Guiana, southern Mexico, Peru, and

northern Argentina. Seeds from Brazil were sent to Florida and the Philippines in the early 1900s. The first written record of rollinia in Hawai'i was not until the mid 1930s, although it may have existed earlier on the homesteads of Filipino immigrants. It is now found growing in most tropical locations and is rapidly becoming a favorite with tropical fruit aficionados.

Cultivars

Currently there are no known breeding programs, although the fruit offers opportunities for breeding and selection. A few known cultivars include 'Regard' in the Philippines, 'Prolific' in Florida, and 'Liso' in Brazil. Fruits of a type called biriba do Alto Solimoes, developed by the Indians living along the Solimoes river in Brazil, can weigh as much as 9 pounds.

Other fruiting *Rollinia* species found mostly in Brazil include *R. sericea*, *R. sylvatica*, *R. salicifolia*, *R. rugulosa*, and *R. emarginata*.

Published by the College of Tropical Agriculture and Human Resources (CTAHR) and issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, under the Director/Dean, Cooperative Extension Service/CTAHR, University of Hawai'i at Mānoa, Honolulu, Hawai'i 96822. Copyright 2010, University of Hawai'i. For reproduction and use permission, contact the CTAHR Office of Communication Services, cos@ctahr.hawaii.edu, 808-956-7036. Hawai'i 96822. an equal opportunity/affirmative action institution providing programs and services to the people of Hawai'i without regard to race, sex, gener identity and expression, age, religion, color, national origin, ancestry, disability, marital status, arrest and court record, sexual orientation, or status as a covered veteran. Find CTAHR publications at www.ctahr.hawaii.edu/freepubs.

Environment

This strictly tropical tree needs warm, most soils in hot, humid climates. It thrives with regular year-round irrigation or in areas where rainfall is equally distributed throughout the year. It does not tolerate cold.

The tree grows best in deep, rich, organic soil and benefits form copious amounts of mulch as long as there is good drainage, but it tolerates poorer and highly acidic soils as long as there is sufficient water.

Propagation

Rollinia is often propagated from seed, which should be planted as soon as possible after harvest. Preferably take seeds from trees that bear regularly with superior fruit. Germination takes about a month and averages 80 percent success. Air-layering and grafting are possible.

Culture and management

Water is the most crucial element when growing rollina for home or commercial production. A tree at 400-foot elevation was irrigated 15 minutes per day from a 1 gallon per hour bubbler emitter, this produced good softball sized or larger fruit year around. The tree remained healthy and flushed until the irrigation was turned off and the tree went into rapid dieback with multiple dead branches. In three months, the unirrigated tree was barely surviving until irrigation was restored, dead wood was removed and NPK 6-6-6 applied. The tree was again thriving within 2 months. The rapid growing tree needs to be pruned regularly to facilitate harvesting as well as for removal of dead wood. It's also advisable to trim extremely long branches as the weight of multiple fruits can cause the branch to break. There are no nutritional studies specifically for Rollina but growers in Brazil as well as Hawai'i usually follow guidelines established for cherimoya, quarterly applications of NPK 6-6-6 or NPK 10-10-10 of up to three pounds per year for mature trees. It is advisable to have a soil analysis to determine needs of micronutrients.

Pests and diseases

There are many problems that could potentially affect commercial Rollinia growing in Hawai'i although few have been observed on healthy well cared for trees.

Moth larva (*Lepidopterene*) can attack maturing fruit. A borer (*Cratosomus bombina*) burrows into the bark and trunk causing branch death. Once pruned, these branches should be burned or disposed of and not left



Rollinia flowers

in the field. White flies (*Aleurodyscus cocois*) and mealy bugs (*Pseudococcus brevipes* and *Aspidiotus destructor*) are found on leaves. *Cercospora* leaf spot occurs and *Glomerella cingulata* causes stem dieback and fruit rot.

Harvesting and yield

Trees at lower elevations in Hawai'i can be found with flowers and all stages of fruit the year around. At higher elevations, between 1000 and 3000 feet, the tree usually sets fruit in the fall, depending on rainfall and irrigation. In Brazil, rollinia trees average 35 to 55 fruits per year but can bear as many as 150 fruits per year on 15-year old trees. South Kona trees produced from 25 to 100 fruits per year.

Commercial fruit is generally harvested when it starts to soften and turn yellow. Care in handling is highly recommended, as the fruit protuberances and skin will blacken considerably when touched.

Postharvest considerations

There are no formal postharvest studies but observations in Hawai'i indicate that once harvested, the fruit will fully ripen in 5 to 8 days. As with many other *Annonaceae*, the fruit will have a short, less than 1 week, shelf life. Once the fruit skin turns fully black, the pulp color and viscosity changes to a clear mucous hence the botanic name, *R. mucosa*. At this stage, the fruit remains edible for only 1 or 2 days before fermentation begins.

Packaging, pricing, and marketing

Rollinia should be harvested and packaged with great care. Hawai'i growers usually use copious amounts of newspaper or bubble wrap to cushion the fruit for transport. Sold at farmers markets, health food stores and small groceries, Rollinia's rapid increase in popularity and high value has made it highly sought after by the store produce managers.

Smaller fruit are sometimes wrapped and sold in Styrofoam netting.

At all markets, fruit is either sold by size or by weight, from \$1.00 for a small fruit to upwards of \$7.00 for an extra large size. This is about twice the price of cherimoya and soursop. By weight, the fruit wholesales for \$3.50 to \$5.00 per pound. Chefs are just beginning to work with the fruit by creating a myriad of sauces, ice creams, flans and other culinary creations. They are clamouring for more fruit. Clearly rollina has a good potential for Hawai'i growers.

Nutritive value

Composition of edible flesh per 100 g

77.2	
80	
2.8	
0.2	
19.1	
1.3	
0.7	
24	
1.2	
26	
33	
0.5	
	77.2 80 2.8 0.2 19.1 1.3 0.7 24 1.2 26 33 0.5

Culinary and other uses

The fruit, often described as having a caramel or lemoncustard-pudding flavor, is usually eaten out of hand. It is often juiced in Brazil and sometimes blended with milk for a drink. It has also been made into wine.



Rollina Soufflé Marg Love

14 oz 1 tsp 1 c	rollina puree coconut extract heavy cream
⁷ / ₂ c 2 tbsp 4	sifted flour extra large eggs, room temperature, separated
2 tbsp	granulated sugar

Reduce 14 oz of rollina puree in a heavy saucepan over low heat and stirring constantly until it is reduced to 10 oz. Remove from heat, stir in coconut extract, and set aside in a cool place. Preheat oven to 350 degrees and lightly butter bottom and sides of soufflé dish. Place 1 cup of heavy cream in saucepan and bring to boil while continually whisking. Remove from heat and set aside. In a medium-sized bowl, mix flour and powdered sugar, then add egg yolks and mix until thick. Drizzle in hot cream, then add mixture back into the same saucepan and cook until it just thickens (less than 5 minutes). Once thickened, place mixture into a bowl and add the rollinia puree. In a separate bowl, beat egg whites to soft peaks, then add granulated sugar and beat until peaks are firm. Gently fold egg whites into rollina mixture, then pour into a soufflé dish. Place filled soufflé dish inside a larger pan filled with hot water that reaches up to half the sides of the soufflé dish. Bake on the middle shelf of oven for 40 minutes at 350 degrees. Carefully remove and serve immediately.

Cost of production

It is essential that growers determine their own cost of production for each crop in each growing location. Including *all* the variables in figuring your cost to produce a specific crop is key to farm sustainability. A few of the operating (or "variable") costs include fertilizer, weed control, pest control, pruning, irrigation, harvesting, marketing, and operations overhead. Ownership (or "fixed") costs also need to be taken into account. For detailed information on the various types of cost, see "The economics of cacao production in Kona" (www. ctahr.hawaii.edu/oc/freepubs/pdf/AB-17.pdf).

The cost-of-production spreadsheet on the following pages can be downloaded as a Microsoft Excel file from www.ctahr.hawaii.edu/oc/freepubs/spreads/6fruits.xls.

Selected references

- Janick, Jules, and Robert E. Paull 2008. The encyclopedia of fruits and nuts. CABI Wallingford, Oxon, UK. p. 68–70.
- Kennard, William C., and Harold F. Winters. 1960. Some fruits and nuts of the tropics. USDA ARS, Misc. Pub. 801.
- Lorenzi, Harri. 1998. Brazilian trees (vol. 2). Instituto Plantarum de Estudos da Flora Ltda. p. 31–32.
- Lorenzi, Harri, et al. 2006. Brazilian fruits and cultivated exotics. Instituto Plantarum de Estudos da Flora Ltda. p. 52–57.
- Silva, Silvestre, and Helena Tassara. 2003. Frutas no Brasil. Editara Editora Ltda. Brazil [in Portuguese]. p. 54–55.

Internet resources

Fruits of warm climates, by Julia F. Morton www.hort.purdue.edu/newcrop/morton/index.html Montoso Gardens www.montosogardens.com Plant Resources of Southeast Asia www.prosea.lipi.go.id International Tropical Fruit Network www.itfnet.org

Acknowledgments

We thank Dr. Francis Zee, Dr. Bryan Brunner, and Gerry Herbert for review and comment on the manuscript, and Dale Evans for editing and layout services. The provision of cost-of-production tables by Dr. Kent Fleming is gratefully acknowledged. UH-CTAHR

Assumptions: (Data entries are annual amounts expressed on a				ssed on a per	tree basis)	Fruit tree =>	ROLLINIA		
1.	Avera	age number of bearing trees (coun	ted)	2	trees	To calculate profi	tability: Enter wage ra	te & benefits actually pai	d (or the rates
2.	Yield	(expressed in number of fruit [F] or lbs)	Р	45	lbs. / tree	one would to pay	r if labor were hired.) T	o calculate <u>cash flow</u> ent	er nothing.
3.	Avera	age wt. (ozs.) / fruit =	20.0	ounces		The cash flow res	ult is (except for deprecia	tion considerations) one's ta	xable income.
4.	Total	lbs. harvested/ tree =	45.0	lbs. gross yi	ield	6.	Wage rate (\$/hr.)	=	\$12.00
5.	Mark	etable yield /tree (%) =	100%	of the gross	s yield	7.	Benefits (FICA, e	etc.) (%) =	33.3%
Gri	ana D	0.000.001	% of tota	¢//b.	I ho /troo/ur		¢ /troo /urr :	¢ /total area /ur	% of groop
1		evenue:	% OF LOCA	ې(۵۱/م) ۵ مارچ	LDS./tree/yr.	markatahla lha	ې /tree /yr.: 20.25	ې /total crop /yr. ۱۱	% of gross
2	Rotai	Lealee	85%	6.45	38.3	marketable lbs.	20.23	41	07%
2	Total	sales = Weighted ave	$\frac{000}{h} =$	\$5 933	45.0	marketable lbs.	266.96	534	100%
Οn	oratir	na Costs:	Enter uni	t quantitios a	as total per vea	r ner tree:	200.00		
<u>ορ</u> Δ	Grov	ving costs:	Liner uni	s/unit	is iolai per yea	¢ /lb of fruit	\$ /tree /vr ·	\$ /total crop /vr	% of gross
Π,	1 F	ertiliziation	onito.	ψ/umr.	Sub-totals =>	0.09	4.27	8.53	2%
	· 🖻	Fertilizer (lbs.)	2.0	\$0.80		0.036	1.60	3.20	_/*
		Labor (min.)	10	\$0.27		0.059	2.67	5.33	
	2 Ir	rination: Assuming ag water rate =	\$2.00	/1 000 gals	Sub total->	0.03	1 3/	2.67	1%
	2 💾	Water (gallons)	ψ <u>2</u> .00	\$0.0	$\frac{300-101a}{100}$	0.00	0.00	0.00	170
		Labor (min.)	5	\$0.27	<u>02</u> / 1,000 guis.	0.000	1 33	2.67	
	2 F			↓ ₩ 0:.21	Quile totale =>	0.00	0.07	5.00	40/
	3 <u>P</u>	est control: Materiale	0.0	¢0.00	Sub-totals =>	0.00	2.07	0.00	1%
		Materials	0.0	\$0.00 \$0.27		0.000	0.00	0.00	
			10	φ0.27		0.059	2.07	0.00	
	4 <u>V</u>	/eed control:			Sub-totals =>	0.03	1.33	2.67	0%
		Chemicals and/or machinery	0.0	\$0.00		0.000	0.00	0.00	
	_	Labor (min.)	5	\$0.27		0.030	1.33	2.67	
	5 <u>P</u>	runing:		1	Sub-totals =>	0.09	4.00	8.00	1%
		Machinery	0.0	\$0.00		0.000	0.00	0.00	
		Labor (min.)	15	\$0.27		0.089	4.00	8.00	
	6	<u>ther:</u>			Sub-totals =>	0.00	0.00	0.00	0%
		Materials and/or machinery	0.0	0.00		0.000	0.00	0.00	
		Labor (min.)	0	\$0.27		0.000	0.00	0.00	
				Total gro	owing costs =	0.302	13.60	27.21	5%
		Enter picking costs based on gros	s yield ar	nd packing a	nd delivery cos	ts based on m	arketable yield .		
В.	Harv	esting costs: Av	erage ce	nts per pou	nd	¢ /lb. of fruit	\$ /tree /yr.:	\$ /enterprise /yr.	% of gross
	1	Picking	20.0	¢/lb.		20.0	9.00	18.00	3%
	2a	Packing: for wholesale	11.6	¢/lb.		2.0	0.14	0.27	0%
	2h	Packing: for retai sales	12.4	¢/lb		12.0	4 74	9.49	2%
	20	<u>Palivani ta market</u>	12.4	ψ/π). ∡ πь		12.0	5.00	9.49	2/0
	3	Delivery to market	11.1	¢/ID.		11.1	5.00	9.99	2%
			Total harve	esting costs =	43.1	18.74	37.48	7%	
		1	OTAL Oper	ating Costs =	43.4	32.34	64.68	12%	
Break-even analysis:			Gross	Margin =	549.8	234.62	469.24	87.9%	
Given the weighted average price of \$5.933 \$/lb. fruit, the mkt. yield required to cover operating costs =					10.9				
Given the marketable yield of 45.0 lbs. fruit/ tree, the ave. price reg. to cover operating costs = \$0.719						\$0.719			

UH-CT	AHR Rollinia	F_N-21 -	- June 2011				
Diekine	How to calculate your harvesting costs expressed as ϕ / lb:						
PICKING 1	Weigh all of the fruit picked in one harvest year & average it out for one tree. Ave. gross yield / tree =	45.0	lbs./year				
0	(Important: The picked fruit yield recorded here is the gross yield and not the marketable yield.)	45	Trainutan				
Z	(Note: You will probably bayest the tree a number of times during the season. We need the time it takes for the wi	40 volo crop voar)	Ininules				
3	<u>(Note</u> , for white probably haves the tree a number of times during the season, we need the time it takes for the wi	inute =	10				
4	Divide the hourly wage rate for nickers by 60 minutes. This will give you the cents per minute wage rate =	<u>nute</u> –	20.0				
5 Divide this wage rate in t / min (result from step 4 above) by the ave nicking rate (in lbs / min) (from step 3 above)							
	The result is your cost (in ¢ / Ib.) to pick a tree's annual gross yield of fruit	20.0	¢ / Ib.				
Exan	nple to illustarate the process:						
a	r average picking r	rate is:					
b C	You would pay pickers \$12.00 per hour = 20 ¢ per minute to pick fruit. $12 \div 60 = 0.20 oYour picking cost / tree is: $20 ¢/min \div 8$ lbs./ min. = 2.5 ¢/ lb . per pound of fruit picked	r 20¢ per minute					
Packing	<u>g:</u> Namo Fonde De la contra de la contr		7				
1	WHOLESALE: Record the total annual cost for packaging to pack the marketable fruit sold wholesale.	<u>\$0.78</u>					
2	Divide this cost by pounds of fruit sold wholesale. (This has been calculated in "Gross Revenue" above)	0.8 11 6	4 / Ib				
3	If more labor (in addition to the nicking labor) is required to pack calculate its cost in d / lb, as above	11.0	¢/m.				
Ū	Extra labor required (minutes): Packing rate = Ibs / minute Iabor cost =						
4	Add these 2 costs together to obtain the total packing cost per pound of fruit marketed wholesale =	11.6	¢/lb.				
5	RETAIL: Follow the same proceedure (steps 1 to 4 above) to calculate the cost to pack fruit sold retail.						
	Total cost of retail packaging = \$4.75 Retail sales = 38.3 pounds Materials cost =	12.4	¢ / Ib.				
	Extra labor required (minutes): 0 Packing rate = Ibs. / minute Labor cost =						
	Total packing cost per pound of fruit marketed retail =	12.4	¢/lb.				
Exan	nple:						
a	In one year you picked 1,600 pounds of fruit, of which 75% was marketable, that is, 1,200 pounds.						
D	During the year you used 24 boxes (@ \$2 each) to ship 1,200 pounds of truit to the wholesale market.	lb of furrite					
C	Divide the packaging cost (\$46) by the amount of marketable fruit. This will give you the materials $\cos t / \frac{1}{2}$	ID. OF ITUIL.					
h	During the year 60 minutes of packing labor was required (beyond the picking labor) Your average pack	king rate is:					
ŭ	1200 lbs. ÷ 60 min. = 20 lbs. / min .						
е	You would pay packers \$12.00 per hour (= 20 ¢ per minute) to pack fruit. Your annual packing labor cos	st /tree is:					
	20 ¢/min ÷ 20 lbs./ min. = 1.0 ¢/ lb.						
f	Add the annual material cost (step c) and labor cost (step e) to obtain your total packing cost / lb. of man	keted fruit.					
	8 ¢/ lb. + 1 ¢ / lb = 9.0 ¢/ lb. for packing wholesale fruit.						
Deliver	<u>y:</u>		-				
1	Based on your annual records, calucuate your average cost / mile for vehicle & driver to haul boxes:	\$1.00	-				
2	Record the total delivery mileage for one year & estimate a portion to allocate to delivering this crop:	0					
3	Record the total weight of marketable fruit delivered during the year.	45.0	¢/lb				
4 Exan	militipy estimated share of mileage times mileage rate & divide by total weight of deliveries.	11.1	¢/10.				
a	You have 10 trees that yield an average of 1.200 lbs of marketable fruit = 12.000 lbs.						
b	During the year you made 24 deliveries carrying 500 lbs of fruit averaging 20 miles round trip.						
С	The cost for your vehicle and driver's time averages about \$1.00 per mile driven.						
	Note: Oviously, the average delivery cost / lb. of all fruit marketed, unlike the picking and packing costs per pound of fruit, will	vary					
	widely for different growers, depending on their location relative to their markets.						
	480 miles driven @ \$1.00 / mile = \$480 \$480.00 transport cost ÷ 12,000 lbs fruit = \$0.04 = 4.0 ¢ / lb	of fruit delivered					