



Vegetative Propagation of Alahee

Harold Ginoza and Fred D. Rauch, Department of Horticulture

Objective

The purpose of this trial was to determine the effect of treatments with a commercial root-promoting product on the rooting of alahee.

Alahee (*Canthium odoratum* [G. Forster] Seem.) is a native Hawaiian plant which has desirable characteristics for tropical landscapes but is difficult to propagate from cuttings (Bornhorst and Rauch 1994). Little information is available on treatments that might enhance the rooting of this plant.

Methods

Uniform, terminal, 4-inch long cuttings of alahee were taken in February from a planting on the UH-Manoa campus. A commercial root-promoting product (Dip'n Grow), which contains 1.0% IBA and 0.5% NAA, was used. The five treatments (Dip'n Grow at 1:1, 1:5, 1:10, and 1:20 dilutions and an untreated control), with 10 cuttings per treatment, were replicated 10 times in a randomized complete block design. The cuttings were placed in vermiculite in 34 cm x 50 cm metal flats in the UH-Manoa Magoon Shade House under intermittent mist (6 sec/2 min cycle).

Dip'n Grow was diluted with water to give the desired concentrations of the chemical. The quick-dip method of application was employed, in which the basal ends of cuttings were dipped in solution for five seconds (Hartmann et al. 1990).

Rooting was evaluated after five months by determining the rooting percentage and the root quality by the method of ranks described by O'Rourke and Maxon (1948) and verified by Mahlstedt and Lana (1958). An index number was determined for each treatment by dividing the sum of scores for each cutting (5 = heavily rooted, 4 = medium rooted, 3 = lightly rooted, 2 = no roots, and 1 = dead) by the number of cuttings.

Results

There was no significant difference among treatments under the conditions of this experiment in root-

ing percentage and root quality. It appears that some other factor besides auxin concentration is limiting the rooting of alahee cuttings.

Literature cited

- Bornhorst, H.L., and F.D. Rauch. 1994. Native Hawaiian plants for landscaping, conservation, and reforestation. Univ. of Hawaii, HITAHR Res. Ext. Series 142.
- Hartmann, H.T., D.E. Kester, and F.T. Davies, Jr. 1990. Plant propagation: principles and practices. Prentice-Hall, Inc., Englewood Cliffs, New Jersey.
- Mahlstedt, J.P., and E.P. Lana. 1958. Evaluation of the rooting response of cuttings by the method of ranks. Proc. Amer. Soc. Hort. Sci. 71:585-590.
- O'Rourke, F.L., and M.A. Maxon. 1948. Effect of particle size of vermiculite media on the rooting of cuttings. Proc. Amer. Soc. Hort. Sci. 51:654-656.

Table 1. The influence of Dip'n Grow on rooting of alahee terminal cuttings after five months.

Dip'n Grow (dilutions)	IBA (ppm)	Rooting percentage	Root quality ^x
control	—	4	1.3 a ^y
1:20	500	5	1.4 a
1:10	1000	7	1.5 a
1:5	2000	7	1.4 a
1:1	5000	7	1.3 a

^xRooting index: 1 = dead, 5 = heavy rooting.

^yMean separation by Duncan's multiple range test, 5% level.

Mention of a trademark, company, or proprietary name does not constitute an endorsement, guarantee, or warranty by the University of Hawaii Cooperative Extension Service or its employees and does not imply recommendation to the exclusion of other suitable products or companies.

Caution: Pesticide use is governed by state and federal regulations. Read the pesticide label to ensure that the intended use is included on it, and follow all label directions.